

**SBDS / Shwachman-Bodian-Diamond
syndrome Antibody, Rabbit PAb,
Antigen Affinity Purified**

Catalog Number: 15362-T54



Sino Biological
Biological Solution Specialist

GENERAL INFORMATION

Immunogen:	Recombinant Human SBDS / Shwachman-Bodian-Diamond syndrome Protein (Catalog#15362-H07E)
Preparation	Produced in rabbits immunized with purified, recombinant Human SBDS / Shwachman-Bodian-Diamond syndrome (rh SBDS / Shwachman-Bodian-Diamond syndrome; Catalog#15362-H07E; Q9Y3A5; Met1-Glu250). SBDS / Shwachman-Bodian-Diamond syndrome specific IgG was purified by Human SBDS / Shwachman-Bodian-Diamond syndrome affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human SBDS / Shwachman-Bodian-Diamond syndrome
Formulation:	0.2 µm filtered solution in PBS
Storage:	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Alternative Names:	CGI-97,SDS,SWDS

APPLICATIONS

Applications:	WB,ELISA,ICC/IF,IP
----------------------	--------------------

RECOMMENDED CONCENTRATION

ICC/IF	ICC/IF: 1:1000-1:5000
Western Blot	WB: 1:500-1:1000
Immunoprecipitation	IP: 1-2 µL/mg of lysate
ELISA	ELISA: 1:25000-1:50000 This antibody can be used at 1:25000-1:50000 with the appropriate secondary reagents to detect Human SBDS / Shwachman-Bodian-Diamond syndrome.


Please Note: Optimal concentrations/dilutions should be determined by the end user.

SBDS / Shwachman-Bodian-Diamond syndrome Antibody, Rabbit PAb, Antigen Affinity Purified


Catalog Number: 15362-T54



Sino Biological
Biological Solution Specialist



Immunofluorescence staining of SBDS in HeLa cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-human SBDS polyclonal antibody (1:5000) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to cytoplasm and nucleus.




Anti-SBDS rabbit polyclonal antibody at 1:500 dilution
Lane A: HepG2 Whole Cell Lysate

Lysates/proteins at 30 µg per lane.

Secondary
Goat Anti- Rabbit IgG H&L (Dylight 800) at 1/10000 dilution.

Developed using the Odyssey technique.
Performed under reducing conditions.

Predicted band size: 29 kDa
Observed band size: 32 kDa



SBDS was immunoprecipitated using:
Lane A: 0.5 mg HepG2 Whole Cell Lysate
Lane B: 0.5 mg Jurkat Whole Cell Lysate

1 µL anti-SBDS rabbit polyclonal antibody and 60 µg of Immunomagnetic beads Protein G.

Primary antibody:
Anti-SBDS rabbit polyclonal antibody, at 1:500 dilution

Secondary antibody:
Dylight 800-labeled antibody to rabbit IgG (H+L), at 1:5000 dilution

Developed using the odssey technique.
Performed under reducing conditions.

Predicted band size: 29 kDa
Observed band size: 29 kDa