## Human NUP210 Insect Cell Lysate (WB positive control)

Catalog Number: 19796-H09BL



## NUP210 Transfected / Overexpression Cell Lysate Product Information

Expressed Host: Baculovirus-Insect Cells

Products Description: Baculovirus-Insect Cell Human NUP210 transfected / overexpressed for Western blot (WB) positive

control. The whole cell lysate is provided in 1X Sample Buffer (1X modified RIPA buffer+1X SDS loading

buffer)

Sequence information: A DNA sequence encoding the human NUP210 (NP\_079199.2) (Leu1837-His1887) was expressed with

a GST tag at the N-terminus.

Predicted N Terminal: Me

Molecule Mass: The recombinant human NUP210 consists of 279 amino acids and predicts a molecular mass of 31.7

kDa.

Species: Human

## **NUP210 Transfected / Overexpression Cell Lysate Usage Guide**

Preparation Method: Cell lysate was prepared by homogenization in ice-cold modified RIPA Lysis Buffer with cocktail of

protease inhibitors (Sigma). Cell debris was removed by centrifugation. Protein concentration was determined by Bradford assay (Bio-Rad protein assay, Microplate Standard assay). The cell lysate was boiled for 5 min in 1 x SDS loading buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium

dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.

Lysis Buffer: Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1%

SDS, 1% Sodium deoxycholate, 1mM PMSF.

**Quality Control Testing:** 12.5% SDS-PAGE Stained with Coomassie Blue after protein purification.

Stability: Samples are stable for up to twelve months from date of receipt.

**Recommend Usage:**1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube.

2. Re-dissolve the pellet using 200µL pure water and boil for 2-5 min.

3. Store the lyophilized cell lysate at 4°C. After re-dissolution, recommend to aliquot it into smaller

quantities and store at -80°C.

Storage Buffer: 1 X Sample Buffer (1 X modified RIPA buffer+1 X SDS loading buffer).

Storage Instruction: Store at 4°C. After re-dissolution, aliquot and store at -80°C.

Application notes: Western blot (WB): Use at an assay dependent dilution.

Other Applications: Not tested.

Optimal dilutions/concentrations should be determined by the end user.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 
■ Tel:+86-400-890-9989 
■ <a href="http://www.sinobiological.com">http://www.sinobiological.com</a>