
LCP1 Antibody, Rabbit PAb, Antigen Affinity Purified






GENERAL INFORMATION	
Immunogen:	E. coli-derived Human LCP1 fragment
Preparation	Produced in rabbits immunized with E. coli-derived Human LCP1 fragment, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human LCP1
Formulation:	0.2 µm filtered solution in PBS
Storage:	This antibody can be stored at $2^{\circ}-8^{\circ}$ for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20° to -80° . Preservative-Free. Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.
APPLICATIONS	
Applications:	WB,IHC-P
RECOMMENDED CONCENTRATION	
IHC-P	IHC-P: 1:500-1:2000
Western Blot	WB: 1:500-1:2000

Please Note: Optimal concentrations/dilutions should be determined by the end user.


LCP1 Antibody, Rabbit PAb, Antigen Affinity Purified

Catalog Number: 200006-T40






Immunochemical staining of human LCP1 in human tonsil with rabbit polyclonal antibody (1:1000, formalin-fixed paraffin embedded sections).



Immunochemical staining of human LCP1 in human spleen with rabbit polyclonal antibody (1:1000, formalin-fixed paraffin embedded sections).



Anti-LCP1 rabbit polyclonal antibody at 1:500 dilution

Lane A: THP1 Whole Cell Lysate Lane B: K562 Whole Cell Lysate Lane C: Jurkat Whole Cell Lysate Lane D: Daudi Whole Cell Lysate

Lysates/proteins at 30 µg per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique. Performed under reducing conditions.

Predicted band size:70 kDa Observed band size:70 kDa