
TRAF4 Antibody, Rabbit PAb, Antigen Affinity Purified






GENERAL INFORMATION	
Immunogen:	E. coli-derived Human TRAF4 fragment
Preparation	Produced in rabbits immunized with E. coli-derived Human TRAF4 fragment, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human TRAF4
Formulation:	PBS, pH7.0 with 0.03% Proclin300
Storage:	This antibody can be stored at $2^{\circ}\text{C-8}^{\circ}\text{C}$ for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C . Avoid repeated freeze-thaw cycles.
Alternative Names:	TRAF4
APPLICATIONS	
Applications:	WB, ICC/IF
RECOMMENDED CONCENTRATION	
ICC/IF	ICC/IF: 1:100-1:500
Western Blot	WB: 1:500-1:2000

Please Note: Optimal concentrations/dilutions should be determined by the end user.


TRAF4 Antibody, Rabbit PAb, Antigen Affinity Purified

Catalog Number: 200187-T34





Immunofluorescence staining of TRAF4 in Hela cells. Cells were fixed with 4% PFA, permeabilzed with 0.1% Triton X-100 in PBS,blocked with 10% serum, and incubated with rabbit anti-Human TRAF4 polyclonal antibody (dilution ratio 1:200) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to Nucleus.



Anti-TRAF4 rabbit polyclonal antibody at 1:500

Lane A: HeLa Whole Cell Lysate Lane B: Jurkat Whole Cell Lysate Lane C: SKBR-3 Whole Cell Lysate Lane D: 293T Whole Cell Lysate

Lysates/proteins at 30 µg per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique. Performed under reducing conditions.

Predicted band size:54 kDa