

STX4 Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological
Biological Solution Specialist

Catalog Number: 200259-T38

GENERAL INFORMATION

Immunogen:	E. coli-derived Human STX4 fragment
Preparation	Produced in rabbits immunized with E. coli-derived Human STX4 fragment, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human STX4
Formulation:	PBS, pH7.0 with 0.03% Proclin300
Storage:	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Avoid repeated freeze-thaw cycles.
Alternative Names:	STX4

APPLICATIONS

Applications:	WB, ICC/IF, IP
---------------	----------------

RECOMMENDED CONCENTRATION

ICC/IF	ICC/IF: 1:100-1:500
Western Blot	WB: 1:500-1:2000
Immunoprecipitation	IP:1-5μL/mg of lysate


Please Note: Optimal concentrations/dilutions should be determined by the end user.

STX4 Antibody, Rabbit PAb, Antigen Affinity Purified


Catalog Number: 200259-T38



Sino Biological
Biological Solution Specialist



Immunofluorescence staining of STX4 in U251MG cells. Cells were fixed with 4% PFA, blocked with 10% serum, and incubated with rabbit anti-Human STX4 polyclonal antibody (dilution ratio 1:200) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to Cytoplasm and cell membrane.




Anti-STX4 rabbit polyclonal antibody at 1:500 dilution

Lane A: HeLa Whole Cell Lysate
Lane B: 293 Whole Cell Lysate
Lane C: K562 Whole Cell Lysate

Lysates/proteins at 30 µg per lane.
Secondary
Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique.
Performed under reducing conditions.

Predicted band size: 34 kDa
Observed band size: 36 kDa



STX4 was immunoprecipitated using:
Lane A: 0.5 mg K562 Whole Cell Lysate

4 µL anti-STX4 rabbit polyclonal antibody and
60 µg of Immunomagnetic beads Protein A/G.

Primary antibody:
Anti-STX4 rabbit polyclonal antibody, at 1:100 dilution

Secondary antibody:
Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution

Developed using the ECL technique.
Performed under reducing conditions.

Predicted band size: 34 kDa
Observed band size: 36 kDa