

PACSIN3 Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological
Biological Solution Specialist

Catalog Number: 200618-T44

GENERAL INFORMATION

Immunogen:	E. coli-derived Human PACSIN3 fragment
Preparation	Produced in rabbits immunized with E. coli-derived Human PACSIN3 fragment, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human PACSIN3
Formulation:	PBS, pH7.0 with 0.03% Proclin300
Storage:	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Avoid repeated freeze-thaw cycles.
Alternative Names:	PACSIN3

APPLICATIONS

Applications:	WB, IHC-P, IP
----------------------	---------------

RECOMMENDED CONCENTRATION

IHC-P	IHC-P: 1:50-1:200
Western Blot	WB: 1:1000-1:5000
Immunoprecipitation	IP: 1-5µL/mg of lysate


Please Note: Optimal concentrations/dilutions should be determined by the end user.

PACSIN3 Antibody, Rabbit PAb, Antigen Affinity Purified


Catalog Number: 200618-T44



Sino Biological
Biological Solution Specialist



Immunohistochemical staining of human PACSIN3 in human placenta with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



Anti-PACSIN3 rabbit polyclonal antibody at 1:500 dilution


Lane A: HepG2 Whole Cell Lysate
Lane B: HeLa Whole Cell Lysate

Lysates/proteins at 30 µg per lane.

Secondary
Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique.
Performed under reducing conditions.

Predicted band size: 48 kDa
Observed band size: 54 kDa



PACSIN3 was immunoprecipitated using:
Lane A: 0.5 mg HepG2 Whole Cell Lysate

4 µL anti-PACSIN3 rabbit polyclonal antibody and 60 µg of Immunomagnetic beads Protein A/G.

Primary antibody:
Anti-PACSIN3 rabbit polyclonal antibody, at 1:100 dilution

Secondary antibody:
Clean-Blot IP Detection Reagent (HRP) at 1:1000 dilution

Developed using the ECL technique.
Performed under reducing conditions.

Predicted band size: 48 kDa
Observed band size: 54 kDa