

MNAT1 Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological
Biological Solution Specialist

Catalog Number: 201154-T38

GENERAL INFORMATION

Immunogen:	E. coli-derived Human MNAT1 fragment
Preparation	Produced in rabbits immunized with E. coli-derived Human MNAT1 fragment, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human MNAT1
Formulation:	PBS, pH7.0 with 0.03% Proclin300
Storage:	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Avoid repeated freeze-thaw cycles.
Alternative Names:	MNAT1
APPLICATIONS	
Applications:	WB, ICC/IF, IP
RECOMMENDED CONCENTRATION	
ICC/IF	ICC/IF: 1:100-1:500
Western Blot	WB: 1:500-1:2000
Immunoprecipitation	IP: 1-5µL/mg of lysate

Please Note: Optimal concentrations/dilutions should be determined by the end user.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

Tel: +86-400-890-9989 (Global), +1-215-583-7898 (USA), +49(0)6196 9678656 (Europe)


Website: <http://www.sinobiological.com>

MNAT1 Antibody, Rabbit PAb, Antigen Affinity Purified




Sino Biological
Biological Solution Specialist

Catalog Number: 201154-T38



Immunofluorescence staining of MNAT1 in A431 cells. Cells were fixed with 4% PFA, permeabilized with 0.1% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human MNAT1 polyclonal antibody (dilution ratio 1:200) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to Nucleus.




Anti-MNAT1 rabbit polyclonal antibody at 1:500 dilution
Lane A: HeLa Whole Cell Lysate
Lane B: A431 Whole Cell Lysate
Lane C: 293T Whole Cell Lysate
Lane D: HepG2 Whole Cell Lysate

Lysates/proteins at 30 µg per lane.
Secondary
Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique.
Performed under reducing conditions.

Predicted band size: 36 kDa
Observed band size: 34 kDa



MNAT1 was immunoprecipitated using:
Lane A: 0.5 mg HeLa Whole Cell Lysate

4 µL anti-MNAT1 rabbit polyclonal antibody and 60 µg of Immunomagnetic beads Protein A/G.

Primary antibody:
Anti-MNAT1 rabbit polyclonal antibody, at 1:100 dilution

Secondary antibody:
Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution

Developed using the ECL technique.
Performed under reducing conditions.

Predicted band size: 34 kDa
Observed band size: 35 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

Tel: +86-400-890-9989 (Global), +1-215-583-7898 (USA), +49(0)6196 9678656 (Europe)

Website: <http://www.sinobiological.com>