

PHB/Prohibitin Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological
Biological Solution Specialist

Catalog Number: 201255-T46

GENERAL INFORMATION

Immunogen:	E. coli-derived Human PHB/Prohibitin fragment
Preparation	Produced in rabbits immunized with E. coli-derived Human PHB/Prohibitin fragment, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human PHB/Prohibitin
Formulation:	PBS, pH7.0 with 0.03% Proclin300
Storage:	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Avoid repeated freeze-thaw cycles.
Alternative Names:	PHB

APPLICATIONS

Applications:	WB, IHC-P, ICC/IF, IP
---------------	-----------------------

RECOMMENDED CONCENTRATION

IHC-P	IHC-P: 1:50-1:200
ICC/IF	ICC/IF: 1:50-1:200
Western Blot	WB: 1:500-1:2000
Immunoprecipitation	IP: 1-5µL/mg of lysate


Please Note: Optimal concentrations/dilutions should be determined by the end user.

PHB/Prohibitin Antibody, Rabbit PAb, Antigen Affinity Purified




Sino Biological
Biological Solution Specialist


Catalog Number: 201255-T46



Immunohistochemical staining of human PHB in human kidney with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



Immunofluorescence staining of PHB in A431 cells. Cells were fixed with 4% PFA, permeabilized with 0.1% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human PHB polyclonal antibody (dilution ratio 1:100) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to Cytoplasm.




Anti-PHB rabbit polyclonal antibody at 1:500 dilution

Lane A: HEK-293 Whole Cell Lysate
Lane B: Jurkat Whole Cell Lysate
Lane C: HepG2 Whole Cell Lysate
Lane D: U-87 MG Whole Cell Lysate

Lysates/proteins at 30 µg per lane.
Secondary
Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique.
Performed under reducing conditions.

Predicted band size: 30 kDa



PHB was immunoprecipitated using:
Lane A: 0.5 mg HEK-293 Whole Cell Lysate

4 µL anti-PHB rabbit polyclonal antibody and
60 µg of Immunomagnetic beads Protein A/G.

Primary antibody:
Anti-PHB rabbit polyclonal antibody, at 1:100 dilution

Secondary antibody:
Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution

Developed using the ECL technique.
Performed under reducing conditions.

Predicted band size: 27 kDa
Observed band size: 27 kDa