

# UQCRC2 Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological  
Biological Solution Specialist

Catalog Number: 201457-T10

GENERAL INFORMATION	
Immunogen:	E. coli-derived Human UQCRC2 fragment
Preparation	Produced in rabbits immunized with E. coli-derived Human UQCRC2 fragment, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human, Cynomolgus
	Reacts with Cynomolgus
Formulation:	PBS, pH7.0 with 0.03% Proclin300
Storage:	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Alternative Names:	MC3DN5,QCR2,UQCR2
APPLICATIONS	
Applications:	IHC-P,ICC/IF
RECOMMENDED CONCENTRATION	
IHC-P	IHC-P: 1:50-1:200
ICC/IF	ICC/IF: 1:50-1:200


**Please Note: Optimal concentrations/dilutions should be determined by the end user.**

# UQCRC2 Antibody, Rabbit PAb, Antigen Affinity Purified


Catalog Number: 201457-T10




Sino Biological  
Biological Solution Specialist



Immunochemical staining of human UQCRC2 in human kidney with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



Immunochemical staining of human UQCRC2 in cynomolgus heart with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



Immunofluorescence staining of UQCRC2 in U2OS cells. Cells were fixed with 4% PFA, permeabilized with 0.1% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human UQCRC2 polyclonal antibody (dilution ratio 1:100) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to Cytoplasm.