

Periostin/POSTN Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological
Biological Solution Specialist

Catalog Number: 201466-T10

GENERAL INFORMATION

Immunogen:	E. coli-derived Human Periostin/POSTN fragment
Preparation	Produced in rabbits immunized with E. coli-derived Human Periostin/POSTN fragment, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human Periostin/POSTN
Formulation:	PBS, pH7.0 with 0.03% Proclin300
Storage:	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Alternative Names:	OSF-2,OSF2,PDLPOSTN,periostin,PN

APPLICATIONS

Applications:	IHC-P,ICC/IF
---------------	--------------

RECOMMENDED CONCENTRATION

IHC-P	IHC-P: 1:200-1:1000
ICC/IF	ICC/IF: 1:100-1:500


Please Note: Optimal concentrations/dilutions should be determined by the end user.

Periostin/POSTN Antibody, Rabbit PAb, Antigen Affinity Purified




Sino Biological
Biological Solution Specialist


Catalog Number: 201466-T10



Immunohistochemical staining of human POSTN in human skin with rabbit polyclonal antibody at 1:500 dilution, formalin-fixed paraffin embedded sections.



Immunohistochemical staining of human POSTN in human breast with rabbit polyclonal antibody at 1:500 dilution, formalin-fixed paraffin embedded sections.



Immunofluorescence staining of POSTN in HeLa cells. Cells were fixed with 4% PFA, blocked with 10% serum, and incubated with rabbit anti-Human POSTN polyclonal antibody (dilution ratio 1:200) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to Cytoplasm.