
## NMRAL1 Antibody, Rabbit PAb, Antigen Affinity Purified






GENERAL INFORMATION	
Immunogen:	E. coli-derived Human NMRAL1 fragment
Preparation	Produced in rabbits immunized with E. coli-derived Human NMRAL1 fragment, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human NMRAL1
Formulation:	PBS, pH7.0 with 0.03% Proclin300
Storage:	This antibody can be stored at $2^{\circ}C-8^{\circ}C$ for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at $-20^{\circ}C$ to $-80^{\circ}C$ . Avoid repeated freeze-thaw cycles.
Alternative Names:	NMRAL1
APPLICATIONS	
Applications:	WB, IHC-P, ICC/IF
RECOMMENDED CONCENTRATION	
IHC-P	IHC-P: 1:50-1:200
ICC/IF	ICC/IF: 1:100-1:500
Western Blot	WB: 1:500-1:2000

Please Note: Optimal concentrations/dilutions should be determined by the end user.


## NMRAL1 Antibody, Rabbit PAb, Antigen Affinity Purified

Catalog Number: 201492-T42






Immunochemical staining of human NMRAL1 in human kidney with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



Immunofluorescence staining of NMRAL1 in MCF7 cells. Cells were fixed with 4% PFA, permeabilzed with 0.1% Triton X-100 in PBS,blocked with 10% serum, and incubated with rabbit anti-Human NMRAL1 polyclonal antibody (dilution ratio 1:200) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to Nucleus.



Anti-NMRAL1 rabbit polyclonal antibody at 1:500 dilution
Lane A: Hela Whole Cell Lysate
Lane B: U-251 MG Whole Cell Lysate

Lysates/proteins at 30 µg per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique. Performed under reducing conditions.

Predicted band size:33 kDa Observed band size:33 kDa (We are unsure as to the identity of these extra bands.)