

NANS Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological
Biological Solution Specialist

Catalog Number: 202003-T46

GENERAL INFORMATION

Immunogen:	E. coli-derived Human NANS fragment
Preparation	Produced in rabbits immunized with E. coli-derived Human NANS fragment, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human NANS
Formulation:	PBS, pH7.0 with 0.03% Proclin300
Storage:	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Avoid repeated freeze-thaw cycles.
Alternative Names:	NANS

APPLICATIONS

Applications:	WB, IHC-P, ICC/IF, IP
----------------------	-----------------------

RECOMMENDED CONCENTRATION

IHC-P	IHC-P: 1:50-1:200
ICC/IF	ICC/IF: 1:100-1:500
Western Blot	WB: 1:500-1:2000
Immunoprecipitation	IP: 1-5µL/mg of lysate


Please Note: Optimal concentrations/dilutions should be determined by the end user.

NANS Antibody, Rabbit PAb, Antigen Affinity Purified


Catalog Number: 202003-T46




Sino Biological
Biological Solution Specialist




Immunohistochemical staining of human NANS in human colon carcinoma with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



Immunohistochemical staining of human NANS in human pancreas with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



Immunofluorescence staining of NANS in U2OS cells. Cells were fixed with 4% PFA, permeabilized with 0.1% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human NANS polyclonal antibody (dilution ratio 1:200) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to Nucleus.



Anti-NANS rabbit polyclonal antibody at 1:500 dilution

Lane A: HepG2 Whole Cell Lysate
Lane B: U-251 MG Whole Cell Lysate


Lysates/proteins at 30 µg per lane.

Secondary

Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique.
Performed under reducing conditions.

Predicted band size: 40 kDa
Observed band size: 40 kDa



NANS was immunoprecipitated using:

Lane A: 0.5 mg U-251 MG Whole Cell Lysate

4 µL anti-NANS rabbit polyclonal antibody and
60 µg of Immunomagnetic beads Protein A/G.

Primary antibody:

Anti-NANS rabbit polyclonal antibody, at 1:100 dilution

Secondary antibody:

Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution

Developed using the ECL technique.
Performed under reducing conditions.

Predicted band size: 40 kDa
Observed band size: 40 kDa