
MTRF1L Antibody, Rabbit PAb, Antigen Affinity Purified






GENERAL INFORMATION	
Immunogen:	E. coli-derived Human MTRF1L fragment
Preparation	Produced in rabbits immunized with E. coli-derived Human MTRF1L fragment, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human MTRF1L
Formulation:	PBS, pH7.0 with 0.03% Proclin300
Storage:	This antibody can be stored at $2^{\circ}C-8^{\circ}C$ for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at $-20^{\circ}C$ to $-80^{\circ}C$. Preservative-Free. Avoid repeated freeze-thaw cycles.
Alternative Names:	HMRF1L,MRF1L,mtRF1a,RP1-101K10.1
APPLICATIONS	
Applications:	WB,ICC/IF,IP
RECOMMENDED CONCENTRATION	
ICC/IF	ICC/IF: 1:100-1:500
Western Blot	WB: 1:500-1:2000
Immunoprecipitation	IP:5-10 μL/mg of lysate

Please Note: Optimal concentrations/dilutions should be determined by the end user.


MTRF1L Antibody, Rabbit PAb, Antigen Affinity Purified

Catalog Number: 202100-T38





Immunofluorescence staining of MTRF1L in HT29 cells. Cells were fixed with 4% PFA, permeabilzed with 0.1% Triton X-100 in PBS,blocked with 10% serum, and incubated with rabbit anti-Human MTRF1L polyclonal antibody (dilution ratio 1:200) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue).Positive staining was localized to Nucleus and Cytoplasm.



Anti-MTRF1L rabbit polyclonal antibody at 1:500 dilution Lane A: U-251 MG Whole Cell Lysate


Lysates/proteins at 30 µg per lane.

Secondary

Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique.Performed under reducing conditions.

Predicted band size:43 kDa Observed band size:43 kDa



MTRF1L was immunoprecipitated using: Lane A:0.5 mg U-251 MG Whole Cell Lysate

4 µL anti-MTRF1L rabbit polyclonal antibody and 60 µg of Immunomagnetic beads Protein A/G.Primary antibody:

Anti-MTRF1L rabbit polyclonal antibody,at 1:100 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10001 dilution

Developed using the ECL technique. Performed under reducing conditions.

Predicted band size: 43 kDa Observed band size: 43 kDa