

EIF3E Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological
Biological Solution Specialist

Catalog Number: 202557-T38

GENERAL INFORMATION

Immunogen:	E. coli-derived Human EIF3E fragment
Preparation	Produced in rabbits immunized with E. coli-derived Human EIF3E fragment, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human EIF3E
Formulation:	PBS, pH7.0 with 0.03% Proclin300
Storage:	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Alternative Names:	eIF3-p46, EIF3-P48, EIF3S6, INT6

APPLICATIONS

Applications:	WB, ICC/IF, IP
---------------	----------------

RECOMMENDED CONCENTRATION

ICC/IF	ICC/IF: 1:100-1:500
Western Blot	WB: 1:500-1:2000
Immunoprecipitation	IP: 1-5µL/mg of lysate


Please Note: Optimal concentrations/dilutions should be determined by the end user.

EIF3E Antibody, Rabbit PAb, Antigen Affinity Purified


Catalog Number: 202557-T38



Sino Biological
Biological Solution Specialist



Immunofluorescence staining of EIF3E in U2OS cells. Cells were fixed with 4% PFA, permeabilized with 0.1% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human EIF3E polyclonal antibody (dilution ratio 1:200) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to Cytoplasm.



Anti-EIF3E rabbit polyclonal antibody at 1:500 dilution


Lane A: Jurkat Whole Cell Lysate
Lane B: A431 Whole Cell Lysate
Lane C: HEK293 Whole Cell Lysate
Lane D: 293T Whole Cell Lysate

Lysates/proteins at 30 µg per lane.

Secondary
Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique. Performed under reducing conditions.

Predicted band size: 52 kDa
Observed band size: 52 kDa



EIF3E was immunoprecipitated using:
Lane A: 0.5 mg Jurkat Whole Cell Lysate

4 µL anti-EIF3E rabbit polyclonal antibody and 60 µg of Immunomagnetic beads Protein A/G. Primary antibody:

Anti-EIF3E rabbit polyclonal antibody, at 1:100 dilution

Secondary antibody:
Clean-Blot IP Detection Reagent (HRP) at 1:1000 dilution

Developed using the ECL technique. Performed under reducing conditions.

Predicted band size: 52 kDa
Observed band size: 52 kDa