
## Adrenodoxin/FDX1 Antibody, Rabbit PAb, Antigen Affinity Purified






| GENERAL INFORMATION       |   |
|---------------------------|---|
| Immunogen:                | E. coli-derived Human Adrenodoxin/FDX1 fragment   |
| Preparation               | Produced in rabbits immunized with E. coli-derived Human Adrenodoxin/FDX1 fragment, and purified by antigen affinity chromatography.  |
| Ig Type:                  | Rabbit IgG  |
| Specificity:              | Human Adrenodoxin/FDX1  |
| Formulation:              | PBS, pH7.0 with 0.03% Proclin300  |
| Storage:                  | This antibody can be stored at $2^{\circ}\text{C}-8^{\circ}\text{C}$ for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at $-20^{\circ}\text{C}$ to $-80^{\circ}\text{C}$ . Preservative-Free. Avoid repeated freeze-thaw cycles. |
| Alternative Names:        | Adrenodoxin,FDX1  |
| APPLICATIONS              |   |
| Applications:             | IHC-P,ICC/IF  |
| RECOMMENDED CONCENTRATION |   |
| IHC-P                     | IHC-P: 1:500-1:2000   |
| ICC/IF                    | ICC/IF: 1:100-1:500   |

Please Note: Optimal concentrations/dilutions should be determined by the end user.


## Adrenodoxin/FDX1 Antibody, Rabbit PAb, Antigen Affinity Purified

Catalog Number: 203052-T10






Immunochemical staining of human FDX1 in human placenta with rabbit polyclonal antibody at 1:1000 dilution, formalin-fixed paraffin embedded sections.



Immunochemical staining of human FDX1 in human adrenal gland with rabbit polyclonal antibody at 1:1000 dilution, formalin-fixed paraffin embedded sections.



Immunofluorescence staining of FDX1 in A431 cells. Cells were fixed with 4% PFA, permeabilzed with 0.1% Triton X-100 in PBS,blocked with 10% serum, and incubated with rabbit anti-Human FDX1 polyclonal antibody (dilution ratio 1:200) at 4°C overnight. Then cells were stained with the Alexa Fluor@488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue).Positive staining was localized to Cytoplasm.