
WDR4 Antibody, Rabbit PAb, Antigen Affinity Purified





GENERAL INFORMATION	
Immunogen:	E. coli-derived Human WDR4 fragment
Preparation	Produced in rabbits immunized with E. coli-derived Human WDR4 fragment, and purified by antigen affinity chromatography.
lg Type:	Rabbit IgG
Specificity:	Human WDR4
Formulation:	PBS, pH7.0 with 0.03% Proclin300
Storage:	This antibody can be stored at $2^{\circ}\text{C-8}^{\circ}\text{C}$ for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C . Avoid repeated freeze-thaw cycles.
APPLICATIONS	
Applications:	WB, IHC-P, IP
RECOMMENDED CONCENTRATION	
IHC-P	IHC-P: 1:50-1:200
Western Blot	WB: 1:500-1:2000
Immunoprecipitation	IP:1-5μL/mg of lysate

Please Note: Optimal concentrations/dilutions should be determined by the end user.


WDR4 Antibody, Rabbit PAb, Antigen Affinity Purified

Catalog Number: 203208-T44





Immunochemical staining of human WDR4 in human liver with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.




Anti-WDR4 rabbit polyclonal antibody at 1:500 dilution

Lane A: HeLa Whole Cell Lysate

Lysates/proteins at 30 µg per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique. Performed under reducing conditions.

Predicted band size:45 kDa Observed band size:45 kDa



WDR4 was immunoprecipitated using: Lane A:0.5 mg HeLa Whole Cell Lysate

4 μ L anti-WDR4 rabbit polyclonal antibody and 60 μ g of Immunomagnetic beads Protein A/G.

Primary antibody:

Anti-WDR4 rabbit polyclonal antibody, at 1:100 dilution

Secondary antibody: Clean-Blot IP Detection Reagent (HRP) at 1:1000dilution

Developed using the ECL technique. Performed under reducing conditions.

Predicted band size: 45 kDa Observed band size: 45 kDa