
PDS5A Antibody, Rabbit PAb, Antigen Affinity Purified






GENERAL INFORMATION	
Immunogen:	E. coli-derived Human PDS5A fragment
Preparation	Produced in rabbits immunized with E. coli-derived Human PDS5A fragment, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human PDS5A
Formulation:	PBS, pH7.0 with 0.03% Proclin300
Storage:	This antibody can be stored at $2^{\circ}C-8^{\circ}C$ for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at $-20^{\circ}C$ to $-80^{\circ}C$. Preservative-Free. Avoid repeated freeze-thaw cycles.
Alternative Names:	PIG54,SCC-112,SCC112
APPLICATIONS	
Applications:	WB,ICC/IF
RECOMMENDED CONCENTRATION	
ICC/IF	ICC/IF: 1:100-1:500
Western Blot	WB: 1:500-1:2000

Please Note: Optimal concentrations/dilutions should be determined by the end user.


PDS5A Antibody, Rabbit PAb, Antigen Affinity Purified

Catalog Number: 203627-T34





Immunofluorescence staining of PDS5A in A431 cells. Cells were fixed with 4% PFA, permeabilzed with 0.1% Triton X-100 in PBS,blocked with 10% serum, and incubated with rabbit anti-Human PDS5A polyclonal antibody (dilution ratio 1:200) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to Nucleus.



Anti-PDS5A rabbit polyclonal antibody at 1:500 dilution

Lane A: U-251 MG Whole Cell Lysate

Lysates/proteins at 30 µg per lane.

Secondary
Goat Anti-Rabbit InG (H

Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique.Performed under reducing conditions.

Predicted band size:150 kDa Observed band size:150 kDa