

MIA3 Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological
Biological Solution Specialist

Catalog Number: 203649-T10

GENERAL INFORMATION

| | |
|--------------------|--|
| Immunogen: | E. coli-derived Human MIA3 fragment |
| Preparation | Produced in rabbits immunized with E. coli-derived Human MIA3 fragment, and purified by antigen affinity chromatography. |
| Ig Type: | Rabbit IgG |
| Specificity: | Human MIA3 |
| Formulation: | PBS, pH7.0 with 0.03% Proclin300 |
| Storage: | This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles. |
| Alternative Names: | UNQ6077,TANGO1,TANGO,D320,ARNT,RP11-378J18.4 |

APPLICATIONS

| | |
|---------------|---------------|
| Applications: | IHC-P, ICC/IF |
|---------------|---------------|

RECOMMENDED CONCENTRATION

| | |
|--------|---------------------|
| IHC-P | IHC-P: 1:50-1:200 |
| ICC/IF | ICC/IF: 1:100-1:500 |


Please Note: Optimal concentrations/dilutions should be determined by the end user.

MIA3 Antibody, Rabbit PAb, Antigen Affinity Purified


Catalog Number: 203649-T10




Sino Biological
Biological Solution Specialist



Immunohistochemical staining of human MIA3 in human brain with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



Immunohistochemical staining of human MIA3 in human rectum with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



Immunofluorescence staining of MIA3 in A549 cells. Cells were fixed with 4% PFA, permeabilized with 0.1% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human MIA3 polyclonal antibody (dilution ratio 1:200) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to Cytoplasm.