

ZNF24 Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological
Biological Solution Specialist

Catalog Number: 203819-T34

GENERAL INFORMATION

| | |
|---------------------------|--|
| Immunogen: | E. coli-derived Human ZNF24 fragment |
| Preparation | Produced in rabbits immunized with E. coli-derived Human ZNF24 fragment, and purified by antigen affinity chromatography. |
| Ig Type: | Rabbit IgG |
| Specificity: | Human ZNF24 |
| Formulation: | PBS, pH7.0 with 0.03% Proclin300 |
| Storage: | This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles. |
| Alternative Names: | KOX17,RSG-A,Zfp191,ZNF191,ZSCAN3 |

APPLICATIONS

| | |
|----------------------|-----------|
| Applications: | WB,ICC/IF |
|----------------------|-----------|

RECOMMENDED CONCENTRATION

| | |
|---------------------|---------------------|
| ICC/IF | ICC/IF: 1:100-1:500 |
| Western Blot | WB: 1:500-1:2000 |


Please Note: Optimal concentrations/dilutions should be determined by the end user.

ZNF24 Antibody, Rabbit PAb, Antigen Affinity Purified




Sino Biological
Biological Solution Specialist

Catalog Number: 203819-T34



Immunofluorescence staining of ZNF24 in U251MG cells. Cells were fixed with 4% PFA, permeabilized with 0.1% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human ZNF24 polyclonal antibody (dilution ratio 1:200) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to Nucleus.



Anti-ZNF24 rabbit polyclonal antibody at 1:500 dilution

Lane A: Jurkat Whole Cell Lysate

Lane B: HeLa Whole Cell Lysate

Lane C: 293 Whole Cell Lysate

Lane D: U-251 MG Whole Cell Lysate

Lysates/proteins at 30 µg per lane.

Secondary
Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique. Performed under reducing conditions.

Predicted band size: 42 kDa
Observed band size: 45 kDa
(We are unsure as to the identity of these extra bands.)