
TRIM33 Antibody, Rabbit PAb, Antigen Affinity Purified






GENERAL INFORMATION	
Immunogen:	E. coli-derived Human TRIM33 fragment
Preparation	Produced in rabbits immunized with E. coli-derived Human TRIM33 fragment, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human TRIM33
Formulation:	PBS, pH7.0 with 0.03% Proclin300
Storage:	This antibody can be stored at $2^{\circ}C-8^{\circ}C$ for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at $-20^{\circ}C$ to $-80^{\circ}C$. Preservative-Free. Avoid repeated freeze-thaw cycles.
Alternative Names:	TRIM33
APPLICATIONS	
Applications:	IHC-P,ICC/IF
RECOMMENDED CONCENTRATION	
IHC-P	IHC-P: 1:500-1:5000
ICC/IF	ICC/IF: 1:100-1:500

Please Note: Optimal concentrations/dilutions should be determined by the end user.


TRIM33 Antibody, Rabbit PAb, Antigen Affinity Purified

Catalog Number: 204438-T10






Immunochemical staining of human TRIM33 in human brain with rabbit polyclonal antibody at 1:2000 dilution, formalin-fixed paraffin embedded sections.



Immunochemical staining of human TRIM33 in human adrenal gland with rabbit polyclonal antibody at 1:2000 dilution, formalin-fixed paraffin embedded sections.



Immunofluorescence staining of TRIM33 in U251MG cells. Cells were fixed with 4% PFA, permeabilzed with 0.1% Triton X-100 in PBS,blocked with 10% serum, and incubated with rabbit anti-Human TRIM33 polyclonal antibody (dilution ratio 1:200) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to .