
ODR4/C1orf27 Antibody, Rabbit PAb, Antigen Affinity Purified






| GENERAL INFORMATION | |
|---------------------------|---|
| Immunogen: | E. coli-derived Human ODR4/C1orf27 fragment |
| Preparation | Produced in rabbits immunized with E. coli-derived Human ODR4/C1orf27 fragment, and purified by antigen affinity chromatography. |
| Ig Type: | Rabbit IgG |
| Specificity: | Human ODR4/C1orf27 |
| Formulation: | PBS, pH7.0 with 0.03% Proclin300 |
| Storage: | This antibody can be stored at $2^{\circ}C-8^{\circ}C$ for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at $-20^{\circ}C$ to $-80^{\circ}C$. Preservative-Free. Avoid repeated freeze-thaw cycles. |
| Alternative Names: | odr-4,ODR4 |
| APPLICATIONS | |
| Applications: | ICC/IF |
| RECOMMENDED CONCENTRATION | |
| ICC/IF | ICC/IF: 1:100-1:500 |

Please Note: Optimal concentrations/dilutions should be determined by the end user.

ODR4/C1orf27 Antibody, Rabbit PAb, Antigen Affinity Purified

Catalog Number: 204562-T02





Immunofluorescence staining of C1orf27 in U2OS cells. Cells were fixed with 4% PFA, permeabilzed with 0.1% Triton X-100 in PBS,blocked with 10% serum, and incubated with rabbit anti-Human C1orf27 polyclonal antibody (dilution ratio 1:200) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue).Positive staining was localized to Cytoplasm and cell membrane.