Influenza A H1N1 (A/Texas/05/2009) Hemagglutinin / HA HEK293 Cell Lysate (WB positive control)

Catalog Number: 40006-V08HL



Hemagglutinin / HA Transfected / Overexpression Cell Lysate Product Information

Expressed Host: HEK293 Cells

Products Description: Human Cell lysate that Influenza A H1N1 (A/Texas/05/2009) HA transfected / overexpressed for Western

blot (WB) positive control. The whole cell lysate is provided in 1X Sample Buffer (1X modified RIPA

buffer+1X SDS loading buffer).

Sequence information: A DNA sequence encoding the extracellular domain of the influenza A H1N1 (A/Texas/05/2009 (H1N1))

hemagglutinin (ACP41934.1) (Met 1-Gln 529, HA1+HA2, uncleved), was fused with a C-terminal

polyhistidine tag.

Predicted N Terminal: Asp 18

Molecule Mass: The secreted recombinant influenza A H1N1 hemagglutinin (A/Texas/05/2009 (H1N1)) comprises 523

amino acids and has a predicted molecular mass of 59 kDa. As a result of glycosylation, the apparent molecular mass of the protein is approximately 70-80 kDa in SDS-PAGE under reducing conditions.

Species: H1N1

Hemagglutinin / HA Transfected / Overexpression Cell Lysate Usage Guide

Preparation Method: Cell lysate was prepared by homogenization in ice-cold modified RIPA Lysis Buffer with cocktail of

protease inhibitors (Sigma). Cell debris was removed by centrifugation. Protein concentration was determined by Bradford assay (Bio-Rad protein assay, Microplate Standard assay). The cell lysate was boiled for 5 min in 1 x SDS loading buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium

dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.

Lysis Buffer: Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1%

SDS, 1% Sodium deoxycholate, 1mM PMSF.

Quality Control Testing: 12.5% SDS-PAGE Stained with Coomassie Blue after protein purification.

Stability: Samples are stable for up to twelve months from date of receipt.

Recommend Usage:1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube.

2. Re-dissolve the pellet using 200µL pure water and boil for 2-5 min.

3. Store the lyophilized cell lysate at 4° C. After re-dissolution, recommend to aliquot it into smaller

quantities and store at -80°C.

Storage Buffer: 1 X Sample Buffer (1 X modified RIPA buffer+1 X SDS loading buffer).

Storage Instruction: Store at 4°C. After re-dissolution, aliquot and store at -80°C.

Application notes: Western blot (WB): Use at an assay dependent dilution.

Other Applications: Not tested.

Optimal dilutions/concentrations should be determined by the end user.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898