

# Dengue virus (DENV) (type 2, strain New Guinea C) NS1 Antibody, Rabbit PAb, Antigen Affinity Purified

Catalog Number: 40263-T62



Sino Biological  
Biological Solution Specialist

## GENERAL INFORMATION

<b>Immunogen:</b>	Recombinant Dengue virus (DENV) (type 2, strain New Guinea C) NS1 Protein (Catalog#40263-V07H)
<b>Preparation</b>	Produced in rabbits immunized with purified, recombinant Dengue virus (DENV) (type 2, strain New Guinea C) NS1 ( Catalog#40263-V07H; AAC59275.1; Asp1-Ala352). Dengue virus (DENV) (type 2, strain New Guinea C) NS1 specific IgG was purified by Dengue virus (DENV) (type 2, strain New Guinea C) NS1 affinity chromatography.
<b>Ig Type:</b>	Rabbit IgG
<b>Specificity:</b>	Dengue virus (DENV) (type 2, strain New Guinea C) NS1
<b>Formulation:</b>	0.2 µm filtered solution in PBS
<b>Storage:</b>	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
<b>Alternative Names:</b>	DENV-NS1

## APPLICATIONS

<b>Applications:</b>	WB,ELISA,IHC-P,FCM,ICC/IF,IP
----------------------	------------------------------

## RECOMMENDED CONCENTRATION

<b>Western Blot</b>	WB: 1:2000-1:5000
<b>ELISA</b>	ELISA: 1:5000-1:10000 This antibody can be used at 1:5000-1:10000 with the appropriate secondary reagents to detect Dengue virus (DENV) (type 2, strain New Guinea C) NS1.


**Please Note: Optimal concentrations/dilutions should be determined by the end user.**

**Dengue virus (DENV) (type 2, strain  
New Guinea C) NS1 Antibody, Rabbit  
PAb, Antigen Affinity Purified**

Catalog Number: 40263-T62



Sino Biological  
Biological Solution Specialist



Anti-Dengue virus (DENV) (type 2, strain New Guinea C) NS1 rabbit polyclonal antibody at 1:2000 dilution

Sample: Dengue virus (DENV) (type 2, strain New Guinea C) NS1 Recombinant Protein

Lane A: 30ng

Lane B: 10ng

Secondary

Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique.  
Performed under reducing conditions.