

Gremlin 1 / GREM1 Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological
Biological Solution Specialist

Catalog Number: 50016-RP02

GENERAL INFORMATION	
Immunogen:	Recombinant Mouse Gremlin 1 / GREM1 protein (Catalog#50016-M08B)
Preparation	Produced in rabbits immunized with purified, recombinant Mouse Gremlin 1 / GREM1 (rM Gremlin 1 / GREM1; Catalog#50016-M08B; O70326; Met1-Asp184). Gremlin 1 / GREM1 specific IgG was purified by Mouse Gremlin 1 / GREM1 affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Mouse Gremlin 1 / GREM1
Formulation:	0.2 µm filtered solution in PBS
Storage:	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.
APPLICATIONS	
Applications:	WB,IHC-P
RECOMMENDED CONCENTRATION	
IHC-P	IHC-P: 1:100-1:500
Western Blot	WB: 1:500-1:2000


Please Note: Optimal concentrations/dilutions should be determined by the end user.

Gremlin 1 / GREM1 Antibody, Rabbit PAb, Antigen Affinity Purified




Sino Biological
Biological Solution Specialist


Catalog Number: 50016-RP02




Immunohistochemical staining of mouse GREM1 in mouse intestine with rabbit polyclonal antibody (1:200, formalin-fixed paraffin embedded sections).



Immunohistochemical staining of mouse GREM1 in mouse heart with rabbit polyclonal antibody (1:200, formalin-fixed paraffin embedded sections).



Immunohistochemical staining of mouse GREM1 in mouse brain with rabbit polyclonal antibody (1:200, formalin-fixed paraffin embedded sections).



Anti-GREM1 rabbit polyclonal antibody at 1:500 dilution
Lane A: HepG2 Whole Cell Lysate

Lysates/proteins at 30 µg per lane.
Secondary
Goat Anti-Rabbit IgG H&L (Dylight800) at 1/10000 dilution.

Developed using the Odyssey technique.
Performed under reducing conditions.

Predicted band size: 21 kDa
Observed band size: 19 kDa