

Mouse IL13RA2 / CD213A2 Protein (His Tag)

Catalog Number: 50061-M08H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

CD213a2

Protein Construction:

A DNA sequence encoding the extracellular domain (Met 1-Lys 334) of mouse IL13R α 2 (NP_032382.1) precursor was fused with a polyhistidine tag at the C-terminus.

Source: Mouse

Expression Host: HEK293 Cells

QC Testing

Purity: > 98 % as determined by SDS-PAGE

Bio Activity:

Measured by its ability to inhibit IL13-dependent proliferation of TF1 human erythroleukemic cells. Kitamura, T. et al. (1989) J. Cell Physiol. 140: 323. The ED₅₀ for this effect is typically 0.1-0.5 μ g/mL.

Endotoxin:

< 1.0 EU per μ g of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Leu 22

Molecular Mass:

The recombinant mouse IL13R α 2 consists of 324 amino acids after removal of the signal peptide and has a predicted molecular mass of 38 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of rIL13R α 2 is approximately 45-50 kDa due to glycosylation.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Interleukin-13 receptor subunit alpha-2 (IL13RA2/IL-13RA2) is also known as also known as cluster of differentiation 213A2 (CD213A2), IL-13 receptor subunit alpha-2, IL-13R subunit alpha-2, and IL-13RA2. The IL13RA2 is often overexpressed in brain tumors, making IL13ra2 one of the vaccine targets for immunotherapy of glioma. IL13RA2/IL-13RA2 is a cancer-associated receptor that is present in greater than 80% of High Grade Astrocytomas (HGA) and has recently been recognized as a cytokine that predisposes breast cancer cells to metastasize. Expression of IL13R α 2 was rapidly lost from the surface of transduced cells grown in culture. The loss appeared to be related to ligands present in fetal bovine serum in the medium. None of the malignant glioma cell lines cultivated in vitro and tested to date exhibited the IL13R α 2 receptor. A recombinant virus (R5111) enters cells via its interaction with the IL13R α 2 receptor in a manner that cannot be differentiated from the interaction of wild-type virus with its receptors.

References

1. Zhou G, *et al.* (2005) Characterization of a recombinant herpes simplex virus 1 designed to enter cells via the IL13R α 2 receptor of malignant glioma cells. *J Virol.* 79(9): 5272-7.
2. Osawa M, *et al.* (2000) Characterization of the mouse interleukin-13 receptor α 1 gene. *Immunogenetics.* 51(11): 974-81.
3. Nair BG, *et al.* (2011) Nanotechnology platforms; an innovative approach to brain tumor therapy. *Med Chem.* 7(5): 488-503.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>