

# Mouse CD27 / TNFRSF7 Protein (His Tag)

Catalog Number: 50110-M08H



Sino Biological  
Biological Solution Specialist

## General Information

### Gene Name Synonym:

S152; Tnfrsf7; Tp55

### Protein Construction:

A DNA sequence encoding the extracellular domain (Met 1-Arg 182) of mouse CD27 (NP\_001028298.1) was fused with a polyhistidine tag at the C-terminus.

**Source:** Mouse

**Expression Host:** HEK293 Cells

## QC Testing

**Purity:** > 96 % as determined by SDS-PAGE

### Bio Activity:

**Immobilized CD70 Protein, Rat, Recombinant (hFc Tag)(Cat:80161-R01H) at 2 µg/mL (100 µL/well) can bind CD27 Protein, Mouse, Recombinant (His Tag)(Cat:50110-M08H), the EC<sub>50</sub> is 50-280 ng/mL.**

### Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

**Predicted N terminal:** Thr 21

### Molecular Mass:

The recombinant mouse CD27 consists of 173 amino acids after removal of the signal peptide and has a predicted molecular mass of 19.6 kDa. In SDS-PAGE under reducing conditions, mCD27 migrates as an approximately 35 kDa band due to glycosylation.

### Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

## Usage Guide

### Stability & Storage:

Samples are stable for twelve months from date of receipt at -20°C to -80°C.


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

**Avoid repeated freeze-thaw cycles.**

### Reconstitution:

Detailed reconstitution instructions are sent along with the products.

## SDS-PAGE:



## Protein Description

CD27, also known as TNFRSF7, is a member of the TNF-receptor superfamily limited to cells of the lymphoid lineage, and exists as both a dimeric glycoprotein on the cell surface and as a soluble protein in serum. As a type I transmembrane glycoprotein of about 55 kDa existing as disulfide-linked homodimer, CD27 has been shown to play roles in lymphoid proliferation, differentiation, and apoptosis. It has an important role in the generation of T cell immunity and is an robust marker for normal memory B cells. It is a T and B cell co-stimulatory molecule, the activity of CD27 is governed by its TNF-like ligand CD70 on lymphocytes and dendritic cells. The CD27-CD70 interaction is required for Th1 generation responses to differentiation signals and long-term maintenance of T cell immunity, and meanwhile, plays a key role in regulating B-cell differentiation, activation and immunoglobulin synthesis.

## References

1. Drner T, *et al.* (2004) Correlation of circulating CD27 high plasma cells and disease activity in systemic lupus erythematosus. *Lupus*. 13(5): 283-9.
2. Sahota SS, *et al.* (2009) CD27 in defining memory B-cell origins in Waldenström's macroglobulinemia. *Clin Lymphoma Myeloma*. 9(1): 33-5.
3. Jiang J, *et al.* (2010) Reduced CD27 expression on antigen-specific CD4+ T cells correlates with persistent active tuberculosis. *J Clin Immunol*. 30(4): 566-73.