

Mouse Oncostatin M / OSM Protein

Catalog Number: 50112-MNAE



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

OncoM

Protein Construction:

A DNA sequence encoding the mouse OSM (P53347) (Ala24-Arg206) was expressed with an initial Met at the N-terminal.

Source: Mouse

Expression Host: E. coli

QC Testing

Purity: > 95 % as determined by SDS-PAGE

Bio Activity:

1. Measured by its ability to inhibit proliferation of M1 mouse myeloid leukemia cells. The ED₅₀ for this effect is typically 10-50 ng/mL. 2. Measured in a cell proliferation assay using NIH3T3 cells. The ED₅₀ for this effect is typically 8-32 ng/mL.

Endotoxin:

Please contact us for more information.

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Met

Molecular Mass:

The recombinant mouse OSM consists of 184 amino acids and predicts a molecular mass of 20.7 KDa. It migrates as an approximately 23 KDa band in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Oncostatin M (OSM) is a glycoprotein belonging to the interleukin-6 family of cytokines that has functions mainly in cell growth. Oncostatin M (OSM) is considered as a pleiotropic cytokine that signals through cell surface receptors type I and type II both of which share the similarity of containing protein gp130 and takes part in many biometabolism processes including liver development, haematopoiesis, inflammation, bone formation and destruction and possibly CNS development. Oncostatin M (OSM) was previously identified by its ability to inhibit the growth of cells from melanoma and other solid tumors. It also has been reported that OSM, like LIF, IL-6 and G-CSF, has the ability to inhibit the proliferation of murine M1 myeloid leukemic cells and can induce their differentiation into macrophage-like cells. The human form of OSM is insensitive between pH2 and 11 and resistant to heating for one hour at 56 degree but is not stable at 9 degrees. The human OSM is produced as a precursor containing 252 amino acids, whose first 25 amino acids function as a secretory signal peptide and which on removal yields the soluble 227 amino acid pro-OSM. Removal of the C-terminal most 31 amino acids produces the fully active 196 residue form.

References

1. Tanaka M, *et al.* (2003) Oncostatin M, a multifunctional cytokine. *Rev Physiol Biochem Pharmacol. Reviews of Physiology, Biochemistry and Pharmacology.* 149: 39-52.
2. Auguste P, *et al.* (1997) Signaling of type II oncostatin M receptor. *J Biol Chem.* 272 (25): 15760-4.
3. Zarlring JM, *et al.* (1986). Oncostatin M: a growth regulator produced by differentiated histiocytic lymphoma cells. *Proc Natl Acad Sci.* 83 (24): 9739-43.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>