

Mouse Thrombopoietin / THPO / TPO Protein (His Tag)

Catalog Number: 50146-M08H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

Mgdf; Ml; Mpllg; Tpo

Protein Construction:

A DNA sequence encoding the mouse THPO (NP_033405.1) precursor (Met 1-Thr 356) with a N-terminal polyhistidine tag was expressed.

Source: Mouse

Expression Host: HEK293 Cells

QC Testing

Purity: > 90 % as determined by SDS-PAGE

Bio Activity:

Measured in a cell proliferation assay using MO7e human megakaryocytic leukemic cells. The ED₅₀ for this effect is typically 1.6-6.4 ng/mL.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Ser 22

Molecular Mass:

The recombinant mouse THPO consists of 346 amino acids with the predicted molecular mass of 37 kDa. As a result of glycosylation, rmTHPO migrates as an approximately 80-90 kDa band in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Thrombopoietin (TPO or THPO), also known as myeloproliferative leukemia virus ligand (c-Mpl), is a hematopoietic growth factor belonging to the EPO/TPO family. The thrombopoietin protein is produced mainly by the liver and the kidney that regulates the production of platelets by the bone marrow. Thrombopoietin protein stimulates both proliferation of progenitor megakaryocytes and their maturation to platelet-producing megakaryocytes, and also accelerates the recovery of platelets. Thrombopoietin protein is involved in cardiovascular disease as it regulates megakaryocyte development and enhances platelet adhesion/aggregation. It has been identified that surface c-MPL, the receptor for thrombopoietin protein, binds to the ligand and mediates the action.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>