

Mouse SLAMF7 / CRACC Protein (His Tag)

Catalog Number: 50201-M08H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

19A; 19A24; 4930560D03Rik; CRACC; CS1

Protein Construction:

A DNA sequence encoding the mouse SLAMF7 (NP_653122.2) extracellular domain (Met 1-Gly 224) was fused with a polyhistidine tag at the C-terminus.

Source: Mouse

Expression Host: HEK293 Cells

QC Testing

Purity: > 95 % as determined by SDS-PAGE

Bio Activity:

Measured by its ability to bind biotinylated mouse SLAMF7 in a functional ELISA.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Ser 23

Molecular Mass:

The recombinant mouse SLAMF7 consists of 213 amino acids with the predicted molecular mass of 23.7 kDa. As a result of glycosylation, the apparent molecular mass of rm SLAMF7 is approximately 35-50 kDa in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

SLAM family member 7 (SLAMF7), also known as CRACC, CD319, CD2-like receptor-activating cytotoxic cells, and CS1, is a single-pass type I membrane protein and a member of the CD2 family of cell surface receptors. SLAMF7 is expressed in NK cells, activated B-cells, NK-cell line but not in promyelocytic, B-cell lines, or T-cell lines. Although the cytoplasmic domain of CS1 contains immunoreceptor tyrosine-based switch motifs (ITSM), which enables to recruit signaling lymphocyte activation molecule (SLAM)-associated protein (SAP/SH2D1A), it activates NK cells in the absence of a functional SAP. CS1 is a self ligand and homophilic interaction of CS1 regulates NK cell cytolytic activity. CRACC positively regulated natural killer cell functions by a mechanism dependent on the adaptor EAT-2 but not the related adaptor SAP. However, in the absence of EAT-2, CRACC potentially inhibited natural killer cell function. It was also inhibitory in T cells, which are typically devoid of EAT-2. Thus, CRACC can exert activating or inhibitory influences on cells of the immune system depending on cellular context and the availability of effector proteins.

References

1. Lee JK, *et al.* (2004) Molecular and functional characterization of a CS1 (CRACC) splice variant expressed in human NK cells that does not contain immunoreceptor tyrosine-based switch motifs. *Eur J Immunol.* 34(10): 2791-9.
2. Tassi I, *et al.* (2005) The cytotoxicity receptor CRACC (CS-1) recruits EAT-2 and activates the PI3K and phospholipase Cgamma signaling pathways in human NK cells. *J Immunol.* 175(12): 7996-8002.
3. Lee JK, *et al.* (2007) CS1 (CRACC, CD319) induces proliferation and autocrine cytokine expression on human B lymphocytes. *J Immunol.* 179(7): 4672-8.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217

• Tel: 215-583-7898

Global Customer: Fax: +86-10-5862-8288

• Tel: +86-400-890-9989

• <http://www.sinobiological.com>