

Mouse CLEC7A / Dectin-1 / CLECSF12 Protein (His Tag)

Catalog Number: 50233-M07H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

beta-GR; BGR; Clecsf12

Protein Construction:

A DNA sequence encoding the extracellular domain (Phe 69-Leu 244) of mouse CLEC7A (NP_064392.2) was fused with a polyhistidine tag at the N-terminus.

Source: Mouse

Expression Host: HEK293 Cells

QC Testing

Purity: > 95 % as determined by SDS-PAGE

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: His

Molecular Mass:

The secreted recombinant mouse CLEC7A consists of 193 amino acids and has a calculated molecular mass of 22.5 kDa. As a result of glycosylation, the apparent molecular mass of the recombinant protein is approximately 30-37 kDa in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Dectin-1 was recently identified as the most important receptor for beta-glucan. It is a type II transmembrane protein which binds beta-1,3 and beta-1,6 glucans, and is expressed on most cells of the innate immune system and has been implicated in phagocytosis as well as killing of fungi by macrophages, neutrophils and dendritic cells. Recognition of beta-glucan by dectin-1 triggers effective immune response, including phagocytosis and proinflammatory factor production, to eliminate infecting fungi, which especially benefits immunocompromised patients against opportunistic fungal infection. In addition, dectin-1 is involved in the adaptive immune response as well as autoimmune diseases and immune tolerance. Dectin-1 can recognize and respond to live fungal pathogens and is being increasingly appreciated as having a key role in the innate responses to these pathogens. In addition to its exogenous ligands, Dectin-1 can recognize an unidentified endogenous ligand on T cells and may act as a co-stimulatory molecule. Recent studies have highlighted the importance of Dectin-1 in anti-fungal immunity, in both mice and humans, and have suggested a possible involvement of this receptor in the control of mycobacterial infections.

References

- 1.Herre J, *et al.* (2004) The role of Dectin-1 in antifungal immunity. *Crit Rev Immunol.* 24(3): 193-203.
- 2.Brown GD. (2006) Dectin-1: a signalling non-TLR pattern-recognition receptor. *6(1):* 33-43.
- 3.Sun L, *et al.* (2007) The biological role of dectin-1 in immune response. *Int Rev Immunol.* 26(5-6): 349-64.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>