

Mouse Cystatin 7 / CST7 Protein (Fc Tag)

Catalog Number: 50236-M02H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

Cmap

Protein Construction:

A DNA sequence encoding the mouse CST7 (O89098) (Met1-Gln144) was expressed, fused with the Fc region of human IgG1 at the C-terminus.

Source: Mouse

Expression Host: HEK293 Cells

QC Testing

Purity: > 93 % as determined by SDS-PAGE

Bio Activity:

Measured by its ability to inhibit active Cathepsin L cleavage of a fluorogenic peptide substrate Z-LR-AMC, R&D Systems, Catalog # ES008. The IC50 is < 6 nM.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Ala 19

Molecular Mass:

The recombinant mouse CST7 /Fc is a disulfide-linked homodimer. The reduced monomer comprises 367 amino acids and has a predicted molecular mass of 41.4 KDa. The apparent molecular mass of the protein is approximately 42 KDa in SDS-PAGE under reducing conditions due to glycosylation.

Formulation:

Lyophilized from sterile 25mM Tris, 0.15M NaCl, pH 7.5

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

The cystatin superfamily members are important natural cysteine protease inhibitors present in a wide variety of organisms and are divided into three classes. Cystatin F, also known as leukocystatin and CMAP (Cystatin-like Metastasis-Associated Protein), is a type 2 cystatin and its expression is limited to hematopoietic cells, with the highest expression levels being observed in monocytes, dendritic cells, and certain types of T-cells. Furthermore, cystatin F mRNA becomes up-regulated during dendritic cell maturation, and thus suggests a specific role of cystatin F in immune regulation. Cystatin F is produced as a dimer, an inactive cathepsin inhibitor which is activated by chemical reduction. In addition, Cystatin F and its homologues have been observed expressing in various human cancer cell lines established from malignant tumors, and thus indicates a new diagnosis and prevention approach of certain human carcinomas metastasis.

References

1. Halfon S., *et al.*, (1998), Leukocystatin, a new class II cystatin expressed selectively by hematopoietic cells. *J. Biol. Chem.* 273:16400-16408.
2. Ni J., *et al.*, (1998), Cystatin F is a glycosylated human low molecular weight cysteine proteinase inhibitor. *J. Biol. Chem.* 273:24797-24804.
3. Morita M., *et al.*, (2000), Genomic construct and mapping of the gene for CMAP (leukocystatin/cystatin F, CST7) and identification of a proximal novel gene, BSCv (C20orf3). *Genomics* 67:87-91.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>