# Mouse IL6RA / CD126 Protein (His Tag)

Catalog Number: 50280-M08H



## **General Information**

### Gene Name Synonym:

IL6R

#### **Protein Construction:**

A DNA sequence encoding the mouse IL6Rα (NP\_034689.2) extracellular domain (Ile 44-Glu 357) was expressed, with a C-terminal polyhistidine tag.

Source: Mouse

Expression Host: HEK293 Cells

**QC** Testing

**Purity:** > 97 % as determined by SDS-PAGE

**Endotoxin:** 

< 1.0 EU per  $\mu g$  of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70  $^{\circ}$ C

Predicted N terminal: Leu 20

## **Molecular Mass:**

The recombinant mouse IL6R $\alpha$  comprises 349 amino acids with a predicted molecular mass of 39 kDa. As a result of glycosylation, it migrates as an approximately 55-60 kDa band in SDS-PAGE under reducing conditions.

#### Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

## **Usage Guide**

#### Storage:


Store it under sterile conditions at  $-20\,^\circ\!\mathrm{C}$  to  $-80\,^\circ\!\mathrm{C}$  upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

### Reconstitution:

Detailed reconstitution instructions are sent along with the products.

#### SDS-PAGE:



# **Protein Description**

Interleukin 6 receptor (IL-6R) also known as CD126 (Cluster of Differentiation 126) is a type I cytokine receptor. The low concentration of a soluble form of IL-6 receptor (sIL-6R) acts as an agonist of IL-6 activity. In the IL-6R/CD126/IL6R system, both a membrane-bound IL-6R and a sIL-6R protein are able to mediate IL-6 signals into the cells through the interaction of gp130. The resulting IL-6/sIL-6R protein complex is also capable of binding to gp130 and inducing intracellular signalling. Through this so-called 'trans-signalling' mechanism, IL-6 is able to stimulate cells that lack an endogenous mIL-6R. High levels of IL-6 and sIL-6R have been reported in several chronic inflammatory and autoimmune diseases as well as in cancer.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898