

# Mouse TGFB3 / Betaglycan Protein (His Tag)

Catalog Number: 50542-M08H



Sino Biological  
Biological Solution Specialist

## General Information

### Gene Name Synonym:

1110036H20Rik; AU015626; AW215636; TBR111

### Protein Construction:

A DNA sequence encoding the extracellular domain of mouse TGFB3 (NP\_072075.2) (Met 1-Arg 399) was expressed, with a polyhistidine tag at the C-terminus.

**Source:** Mouse

**Expression Host:** HEK293 Cells

## QC Testing

**Purity:** > 95 % as determined by SDS-PAGE

### Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

### Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

**Predicted N terminal:** Gly 23

### Molecular Mass:

The recombinant mouse TGFB3 consists of 774 amino acids and has a predicted molecular mass of 86.3 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of rmTGFB3 is approximately 80-90 kDa due to glycosylation.

### Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

## Usage Guide

### Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

**Avoid repeated freeze-thaw cycles.**

### Reconstitution:

Detailed reconstitution instructions are sent along with the products.

## SDS-PAGE:



## Protein Description

Betaglycan also known as transforming growth factor beta receptor III (TGFB3), is a cell-surface chondroitin sulfate / heparan sulfate proteoglycan. TGFB3 is a transforming growth factor (TGF)-beta type III receptor. This receptor is a membrane proteoglycan that often functions as a co-receptor with other TGF-beta receptor superfamily members. Ectodomain shedding produces soluble TGFB3, which may inhibit TGFβ signaling. Decreased expression of this receptor has been observed in various cancers. TGFB3 is the TGF-β component most commonly downregulated among localized human prostate cancer studies. TGFB3 knockdown led to focus formation and enhanced expression of CD133, a marker found on prostate cancer stem cells. TGFB3 is an accessory receptor that binds to and modulates the activities of both transforming growth factor-beta (TGFβ) and inhibin, two members of the TGFβ superfamily of growth factors that regulate many aspects of reproductive biology. TGFB3 is known to be expressed in adult testis and ovary, but little is known about this receptor during gonadogenesis.

## References

1. Johnson DW, *et al.* (1996) Assignment of human transforming growth factor-beta type I and type III receptor genes (TGFB1 and TGFB3) to 9q33-q34 and 1p32-p33, respectively. *Genomics*. 28 (2): 356-7.
2. Rotzer D, *et al.* (2001) Type III TGF-beta receptor-independent signalling of TGF-beta2 via T betaRII-B, an alternatively spliced TGF- type II receptor. *EMBO J*. 20 (3): 480-90.
3. Gao J, *et al.* (1999) Expression of transforming growth factor-beta receptors types II and III within various cells in the rat periodontium. *J Periodont Res*. 34 (2): 113-22.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>