

Mouse CD34 Protein (His Tag)

Catalog Number: 50589-M08H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

AU040960

Protein Construction:

A DNA sequence encoding the extracellular domain of mouse CD34 (NP_001104529.1) (Met 1-Thr 287) was expressed, with a polyhistidine tag at the C-terminus.

Source: Mouse

Expression Host: HEK293 Cells

QC Testing

Purity: > 98 % as determined by SDS-PAGE

Bio Activity:

Measured by the ability of the immobilized protein to support the adhesion of the HUVEC human umbilical vein endothelial cell line . When 4×10^4 cells/well are added to mouse CD34 coated plates (0.8 $\mu\text{g/ml}$, 100 $\mu\text{l/well}$), approximately >30 % will adhere after one hour at 37 $^{\circ}\text{C}$.

Endotoxin:

< 1.0 EU per μg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 $^{\circ}\text{C}$

Predicted N terminal: Thr 35

Molecular Mass:

The recombinant mouse CD34 consists of 264 amino acids and has a predicted molecular mass of 28.6 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of rmCD34 is approximately 50-60 kDa due to glycosylation.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20 $^{\circ}\text{C}$ to -80 $^{\circ}\text{C}$ upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Cluster of Differentiation 34 (CD34) is a member of a family of single-pass transmembrane sialomucin proteins, and may function as a cell-cell adhesion factor. CD34 protein is selectively expressed on hematopoietic progenitor cells and the small vessel endothelium of a variety of tissues. It has been widely used as a stem and progenitor cell marker, and clinical CD34+ stem cell transplantation (CD34+SCT) has been performed for tumor purging. CD34 monoclonal antibodies are widely used to identify and isolate hemopoietic progenitors and to classify acute and chronic leukemias.

References

- 1.Hogan CJ, *et al.* (2002) Differential long-term and multilineage engraftment potential from subfractions of human CD34+ cord blood cells transplanted into NOD/SCID mice. *Proc Nat Acad Sci USA*. 99 (1): 413-8.
- 2.Nielsen JS,*et al.* (2009) CD34 is a key regulator of hematopoietic stem cell trafficking to bone marrow and mast cell progenitor trafficking in the periphery. *Microcirculation*. 16(6): 487-96.
- 3.Mastrandrea F,*et al.* (2009) CD34+ hemopoietic precursor and stem cells traffic in peripheral blood of celiac patients is significantly increased but not directly related to epithelial damage severity. *Eur Ann Allergy Clin Immunol*. 40(3): 90-103.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>