

Mouse PSGL-1 / CD162 / SELPLG Protein (Fc Tag)

Catalog Number: 50770-M02H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

CD162; Psgl-1; Psgl1; Selp1; Selp1

Protein Construction:

A DNA sequence encoding the extracellular domain of mouse SELPLG (Q62170) (Met 1-Cys 307) was fused with the Fc region of human IgG1 at the C-terminus.

Source: Mouse

Expression Host: HEK293 Cells

QC Testing

Purity: > 85 % as determined by SDS-PAGE

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Leu 18

Molecular Mass:

The secreted recombinant mouse SELPLG/Fc is a disulfide-linked homodimer. The reduced monomer comprises 531 amino acids and has a calculated molecular mass of 57.4 kDa. As a result of glycosylation, the apparent molecular mass of rmSELPLG/Fc monomer is approximately 82 kDa in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

P-selectin glycoprotein ligand-1 (PSGL-1), also known as SELPLG or CD162, is the high affinity counter-receptor for P-selectin on expressed on activated endothelial cells and platelets. PSGL-1 is a mucin-type glycoprotein, expressed on leukocytes and platelets as a homodimer of two disulfide-linked subunits of ~120 kD. As cell adhesion molecules, multiple studies have shown that PSGL-1/ P-selectin interaction is required for the normal recruitment of leukocytes during inflammatory reactions, and also participates in hemostatic responses. PSGL-1 protein requires two distinct posttranslational modifications for the Ca²⁺-dependent recognition by the lectin domain of P-selectin, that is tyrosine sulfation and specific O-linked glycosylation (sialic acid and fucose). PSGL-1 can also bind to other two members of the selectin family, E-selectin (endothelial) and L-selectin (leukocyte), but binds best to P-selectin.

References

- 1.1. Sako, D. et al., 1993, Cell. 75: 1179-1186.
- 2.2. Wilkins, P. P. et al., 1995, J. Biol. Chem. 270: 22677-22680.
- 3.3. Frenette, P. S. et al., 2000, J. Exp. Med. 191: 1413-1422.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>