

Mouse TNFRSF4 / OX40 / CD134 Protein (Fc Tag)

Catalog Number: 50808-M02H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

ACT35; CD134; Ly-70; Ox40; Txgp1; TXGP1L

Protein Construction:

A DNA sequence encoding the extracellular domain of mouse TNFRSF4 (NP_035789.1) (Met 1-Pro 211) was fused with the Fc region of human IgG1 at the C-terminus.

Source: Mouse

Expression Host: HEK293 Cells

QC Testing

Purity: > 92 % as determined by SDS-PAGE

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Predicted N terminal: Val 20

Molecular Mass:

The recombinant mouse TNFRSF4/Fc is a disulfide-linked homodimer. The reduced monomer comprises 433 amino acids and has a calculated molecular mass of 48.3 kDa. As a result of glycosylation, the apparent molecular mass of the monomer is approximately 60 kDa in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Stability & Storage:

Samples are stable for twelve months from date of receipt at -20°C to -80°C.


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

OX40 (CD134) and its binding partner, OX40L (CD252), are members of the tumor necrosis factor receptor/tumor necrosis factor superfamily, is known to break an existing state of tolerance in malignancies, leading to a reactivation of antitumor immunity. The interaction between OX40 and OX40L plays an important role in antigen-specific T-cell expansion and survival. OX40 and OX40L also regulate cytokine production from T cells, antigen-presenting cells, natural killer cells, and natural killer T cells, and modulate cytokine receptor signaling. In line with these important modulatory functions, OX40-OX40L interactions have been found to play a central role in the development of multiple inflammatory and autoimmune diseases, making them attractive candidates for intervention in the clinic. Conversely, stimulating OX40 has shown it to be a candidate for therapeutic immunization strategies for cancer and infectious disease.

References

1. Compaan D.M., et al. (2006) .The crystal structure of the costimulatory OX40-OX40L complex. Structure 14:1321-1330.
2. Kawamata S., et al. (1998) .Activation of OX40 signal transduction pathways leads to tumor necrosis factor receptor-associated factor (TRAF) 2- and TRAF5-mediated NF-kappaB activation. J. Biol. Chem. 273:5808-5814.
3. Byun M., (2013) Inherited human OX40 deficiency underlying classic Kaposi sarcoma of childhood. J. Exp. Med. 210:1743-1759.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

Tel: +86-400-890-9989 (Global), +1-215-583-7898 (USA), +49(0)6196 9678656 (Europe)

Website: <http://www.sinobiological.com>