

# Mouse DDR1 Kinase / MCK10 / CD167 Protein (His & GST Tag)

Catalog Number: 50829-M20B1



Sino Biological  
Biological Solution Specialist

## General Information

### Gene Name Synonym:

6030432F18; AI323681; Cak; CD167a; Nep; PTK3A

### Protein Construction:

A DNA sequence encoding the mouse DDR1 (Q03146-2) (Leu444-Val874) was fused with the N-terminal polyhistidine-tagged GST tag at the N-terminus.

**Source:** Mouse

**Expression Host:** Baculovirus-Insect Cells

## QC Testing

**Purity:** > 95 % as determined by SDS-PAGE

### Bio Activity:

The specific activity was determined to be 2 nmol/min/mg using synthetic modified AXLtide peptide (modified-CKKSRGDYMTMQIG) as substrate.

### Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

### Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

**Predicted N terminal:** Met

### Molecular Mass:

The recombinant mouse DDR1/GST chimera consists of 668 amino acids and has a calculated molecular mass of 75.8kDa. The recombinant protein migrates as an approximately 68 kDa band in SDS-PAGE under reducing conditions.

### Formulation:

Supplied as sterile 20mM Tris, 500mM NaCl, pH 7.4, 10% glycerol, 2mM DTT

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

## Usage Guide

### Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

**Avoid repeated freeze-thaw cycles.**

### Reconstitution:

Detailed reconstitution instructions are sent along with the products.

## SDS-PAGE:



## Protein Description

Discoidin domain receptor family, member 1 (DDR1), also known as or CD167a (cluster of differentiation 167a), and Mammary carcinoma kinase 10 (MCK10), belongs to a subfamily of tyrosine kinase receptors with an extracellular domain homologous to Dictyostellium discoideum protein discoidin 1. Receptor tyrosine kinases play a key role in the communication of cells with their microenvironment. These kinases are involved in the regulation of cell growth, differentiation and metabolism. Expression of DDR1/MCK10/CD167 is restricted to epithelial cells, particularly in the kidney, lung, gastrointestinal tract, and brain. In addition, it has been shown to be significantly overexpressed in several human tumors. DDR1/MCK10/CD167 plays an important role in regulating attachment to collagen, chemotaxis, proliferation, and MMP production in smooth muscle cells. DDR1 functions in a feedforward loop to increase p53 levels and at least some of its effectors. Inhibition of DDR1 function resulted in strikingly increased apoptosis of wild-type p53-containing cells in response to genotoxic stress through a caspase-dependent pathway.

## References

- 1.Hou G, *et al.* (2001) The discoidin domain receptor tyrosine kinase DDR1 in arterial wound repair. *J Clin Invest.* 107(6): 727-35.
- 2.Ongusaha PP, *et al.* (2003) p53 induction and activation of DDR1 kinase counteract p53-mediated apoptosis and influence p53 regulation through a positive feedback loop. *EMBO J.* 22(6): 1289-301.
- 3.Jönsson M, *et al.* (2001) Repression of Wnt-5a impairs DDR1 phosphorylation and modifies adhesion and migration of mammary cells. *J Cell Sci.* 114(11): 2043-53.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>