

Mouse Prealbumin / Transthyretin / TTR / PALB Protein (His Tag)

Catalog Number: 50958-M08H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

AA408768; AI787086; D17860; prealbumin

Protein Construction:

A DNA sequence encoding the mouse TTR (NP_038725.1) (Met1-Asn147) was expressed with a polyhistidine tag at the C-terminus.

Source: Mouse

Expression Host: HEK293 Cells

QC Testing

Purity: > (4.2+8.5+6.8+79.1) % as determined by SDS-PAGE.

Endotoxin:

< 1.0 EU per µg protein as determined by the LAL method.

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Gly 21

Molecular Mass:

The recombinant mouse TTR consists of 127 amino acids and predicts a molecular mass of 13.6 kDa.

Formulation:

Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Prealbumin/Transthyretin, also known as ATTR, Prealbumin, TTR and PALB, is a secreted and cytoplasm protein which belongs to the Prealbumin / Transthyretin family. Prealbumin / Transthyretin is detected in serum and cerebrospinal fluid (at protein level). It is highly expressed in choroid plexus epithelial cells. It is also detected in retina pigment epithelium and liver. Each monomer of Prealbumin / Transthyretin has two 4-stranded beta sheets and the shape of a prolate ellipsoid. Antiparallel beta-sheet interactions link monomers into dimers. A short loop from each monomer forms the main dimer-dimer interaction. These two pairs of loops separate the opposed, convex beta-sheets of the dimers to form an internal channel. Prealbumin/Transthyretin is a carrier protein. It transports thyroid hormones in the plasma and cerebrospinal fluid, and also transports retinol (vitamin A) in the plasma. Defects in Prealbumin / Transthyretin are the cause of amyloidosis type 1 (AMY1) which is a hereditary generalized amyloidosis due to Prealbumin / Transthyretin amyloid deposition. Protein fibrils can form in different tissues leading to amyloid polyneuropathies, amyloidotic cardiomyopathy, carpal tunnel syndrome, systemic senile amyloidosis. The diseases caused by mutations include amyloidotic polyneuropathy, euthyroid hyperthyroxinaemia, amyloidotic vitreous opacities, cardiomyopathy, oculoleptomeningeal amyloidosis, meningocerebrovascular amyloidosis, carpal tunnel syndrome, etc.

References

1. Westmark P, *et al.* (1990) Fibril in senile systemic amyloidosis is derived from normal transthyretin. *Proc Natl Acad Sci U S A.* 87(7): 2843-5.
2. Colon W, *et al.* (1992) Partial denaturation of transthyretin is sufficient for amyloid fibril formation in vitro. *Biochemistry.* 31(36): 8654-60.
3. Hammarström P, *et al.* (2003) Prevention of transthyretin amyloid disease by changing protein misfolding energetics. *Science.* 299(5607): 713-6.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217

• Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288

• Tel:+86-400-890-9989

• <http://www.sinobiological.com>