

# Mouse LOXL2 / Lysyl oxidase homolog 2 Protein (His Tag)

Catalog Number: 53068-M08H



Sino Biological  
Biological Solution Specialist

## General Information

### Gene Name Synonym:

1110004B06Rik; 4930526G11Rik; 9430067E15Rik

### Protein Construction:

A DNA sequence encoding the mouse Loxl2 (NP\_201582.2) (Met1-Gln776) was expressed with a polyhistidine tag at the C-terminus.

**Source:** Mouse

**Expression Host:** HEK293 Cells

## QC Testing

**Purity:** > (19.0+61.5)% as determined by SDS-PAGE

### Bio Activity:

**Measured by its ability to produce hydrogen peroxide during the oxidation of benzylamine. The specific activity is > 3pmoles/min/μg.**

### Endotoxin:

< 1.0 EU per μg protein as determined by the LAL method.

### Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

**Predicted N terminal:** Gln 26

### Molecular Mass:

The recombinant mouse Loxl2 consists 762 amino acids and predicts a molecular mass of 85.9 kDa.

### Formulation:

Lyophilized from sterile 20 mM MES, 150 mM NaCl, pH 6.5.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

## Usage Guide

### Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

### Avoid repeated freeze-thaw cycles.

### Reconstitution:

Detailed reconstitution instructions are sent along with the products.

## SDS-PAGE:



## Protein Description

Lysyl oxidase homolog 2, also known as Lysyl oxidase-like protein 2, Lysyl oxidase-related protein 2, Lysyl oxidase-related protein WS9-14 and LOXL2, is a secreted protein which belongs to the lysyl oxidase family. LOXL2 contains four SRCR domains. The lysyl oxidase family is made up of five members: lysyl oxidase (LOX) and lysyl oxidase-like 1-4 ( LOXL1, LOXL2, LOXL3, LOXL4 ). All members share conserved C-terminal catalytic domains that provide for lysyl oxidase or lysyl oxidase-like enzyme activity; and more divergent propeptide regions. LOX family enzyme activities catalyze the final enzymatic conversion required for the formation of normal biosynthetic collagen and elastin cross-links. LOXL2 is expressed by pre-hypertrophic and hypertrophic chondrocytes in vivo, and that LOXL2 expression is regulated in vitro as a function of chondrocyte differentiation. LOXL2 promotes chondrocyte differentiation by mechanisms that are likely to include roles as both a regulator and an effector of chondrocyte differentiation. LOXL2 expression could also be explored as a molecular target in the prevention of breast cancer progression.

## References

- 1.Peng,L. et al., 2009, Carcinogenesis. 30 (10):1660-9.
- 2.Hollosi,P. et al., 2009, Int J Cancer. 125 (2):318-27.
- 3.Rückert,F. et al., 2010, Int J Colorectal Dis. 25 (3):303-11.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>