

# Canine CD40L / CD154 / TNFSF5 Protein



Sino Biological  
Biological Solution Specialist

Catalog Number: 70068-DNCH

## General Information

### Gene Name Synonym:

CD40LG

### Protein Construction:

A DNA sequence encoding the canine CD40LG (O97626) (Met112-Leu260) was expressed with two additional amino acids (Gly & Pro) at the N-terminus.

**Source:** Canine

**Expression Host:** HEK293 Cells

## QC Testing

**Purity:** > 95 % as determined by SDS-PAGE

### Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

### Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

**Predicted N terminal:** Gly

### Molecular Mass:

The recombinant canine CD40LG comprises 151 amino acids and has a predicted molecular mass of 16 kDa. The apparent molecular mass of the protein is approximately 19 kDa in SDS-PAGE under reducing conditions.

### Formulation:

Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

## Usage Guide

### Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

**Avoid repeated freeze-thaw cycles.**

### Reconstitution:

Detailed reconstitution instructions are sent along with the products.

## SDS-PAGE:



## Protein Description

The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. CD154, also known as CD40 ligand or CD40L, is a member of the TNF superfamily. While CD154 was originally found on T cell surface, its expression has since been found on a wide variety of cells, including platelets, mast cells, macrophages and NK cells. CD154's ability is achieved through binding to the CD40 on antigen- presenting cells (APC). In the macrophage cells, the primary signal for activation is IFN-γ from Th1 type CD4 T cells. The secondary signal is CD40L on the T cell, which interacting with the CD40 molecules, helping increase the level of activation.

## References

- 1.Zola H, *et al.* (2007) CD molecules 2006-human cell differentiation molecules. *J Immunol Methods.* 318 (1-2): 1-5.
- 2.Ho IC, *et al.* (2009) GATA3 and the T-cell lineage: essential functions before and after T-helper-2-cell differentiation. *Nat Rev Immunol.* 9 (2): 125-35.
- 3.Matesanz-Isabel J, *et al.* (2011) New B-cell CD molecules. *Immunology Letters.* 134 (2): 104-12.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 ● Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 ● Tel:+86-400-890-9989 ● <http://www.sinobiological.com>