

# Rat CNTFR / CNTFR-alpha Protein (His Tag)



Sino Biological  
Biological Solution Specialist

Catalog Number: 80019-R08B

## General Information

### Gene Name Synonym:

CNTFR

### Protein Construction:

A DNA sequence encoding the rat CNTFR (Q08406-1) (Met 1-Pro 346) was fused with a C-terminal polyhistidine tag.

**Source:** Rat

**Expression Host:** Baculovirus-Insect Cells

## QC Testing

**Purity:** > 97 % as determined by SDS-PAGE

### Bio Activity:

**Measured by its binding ability in a functional ELISA**  
• Immobilized Rat CNTFR at 10 µg/ml (100 µl/well) can bind biotinylated human CNTF with a linear range of 1.28-160 ng/ml

### Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

**Predicted N terminal:** Met 1

### Molecular Mass:

The recombinant rat CNTFR comprises 356 amino acids with a predicted molecular mass of 39.6 kDa. It migrates as an approximately 48 kDa band in reduced SDS-PAGE.

### Formulation:

Lyophilized from sterile 20mM Tris, 500mM NaCl, 10% gly, pH 8.0

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

## Usage Guide

### Stability & Storage:

Samples are stable for twelve months from date of receipt at -20°C to -80°C.


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

**Avoid repeated freeze-thaw cycles.**

### Reconstitution:

Detailed reconstitution instructions are sent along with the products.

## SDS-PAGE:



## Protein Description

Ciliary neurotrophic factor(CNTF) is a member of the cytokine family. It is a polypeptide hormone that have functions in promoting neurotransmitter synthesis and neurite outgrowth in certain neuronal populations. It's actions appear to be restricted to the nervous system. Ciliary neurotrophic factor(CNTF) has biological effects through the activation of a multi-subunit receptor complex, consisting of an extracellular CNTF binding subunit(CNTF&alpha;) and two transmembrane signal transduction proteins: glycoprotein gp130 and LIF receptor. CNTF is considered as a potent survival factor of neurons and oligodendrocytes and may be relevant in reducing tissue destruction during inflammatory attacks. CNTF is also a survival factor for neurons of the peripheral sensory sympathetic, and ciliary ganglia. It has been reported that CNTF could be an agent that has therapeutic potential and possibly induces differentiation of large multipolar ganglionic phenotype in a subset of progenitors.

## References

- 1.Dutt K, et al. (2010) Ciliary neurotrophic factor: a survival and differentiation inducer in human retinal progenitors. In Vitro Cell Dev Biol Anim. 46 (7) : 635-46.
- 2.Lam A, et al. ( 1991) Sequence and structural organization of the human gene encoding ciliary neurotrophic factor. Gene 102 (2) : 271-6.
- 3.Bazan JF. ( 1991) Neuropoietic cytokines in the hematopoietic fold. Neuron 7 (2) : 197-208.