

Rat VEGFC / VEGF-C Protein (aa 108-223, His Tag)

Catalog Number: 80103-R08H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

VEGFC

Protein Construction:

A DNA sequence encoding the rat VEGFC (O35757) (Ala108-Arg223) was expressed, fused with a polyhistidine tag at the C-terminus.

Source: Rat

Expression Host: HEK293 Cells

QC Testing

Purity: > 95 % as determined by SDS-PAGE

Bio Activity:

1. Immobilized rat VEGFC-His at 10 µg/mL (100 µL/well) can bind mouse VEGFR3-Fc (cat:50584-M02H). The EC_{50} of mouse VEGFR3-Fc (cat:50584-M02H) is 17.4-40.6 ng/mL. 2. Measured in a cell proliferation assay using human umbilical vein endothelial cells (HUVEC). The ED_{50} for this effect is typically 0.1-0.8 µg/mL.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Ala 108

Molecular Mass:

The recombinant rat VEGFC comprises 127 amino acids and predicts a molecular mass of 14.5 kDa.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Vascular endothelial growth factor C (VEGF-C) is a member of the VEGF family. Upon biosynthesis, VEGF-C protein is secreted as a non-covalent homodimer in an anti-parallel fashion. VEGF-C protein is a dimeric glycoprotein, as a ligand for two receptors, VEGFR-3 (Flt4), and VEGFR-2. VEGF-C may function in angiogenesis of the venous and lymphatic vascular systems during embryogenesis. VEGF-C protein is over-expressed in various human cancers including breast cancer and prostate cancer. VEGF-C/VEGFR-3 axis, through different signaling pathways, plays a critical role in cancer progression by regulating different cellular functions, such as invasion, proliferation, and resistance to chemotherapy. Thus, targeting the VEGF-C/VEGFR-3 axis may be therapeutically significant for certain types of tumors.

References

1. Joukov V, *et al.* (1997) Vascular endothelial growth factors VEGF-B and VEGF-C. *J Cell Physiol.* 173(2): 211-5.
2. Su JL, *et al.* (2007) The role of the VEGF-C/VEGFR-3 axis in cancer progression. *Br J Cancer.* 96(4): 541-5.
3. Anisimov A, *et al.* (2009) Activated forms of VEGF-C and VEGF-D provide improved vascular function in skeletal muscle. *Circ Res.* 104(11): 1302-12.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217

● Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288

● Tel:+86-400-890-9989 ●

<http://www.sinobiological.com>