

# Rat Ephrin-B1 / EFNB1 HEK293 Cell Lysate (WB positive control)



Sino Biological Inc.

Biological Solution Specialist

Catalog Number: 80106-R08HL

## Ephrin-B1 Transfected / Overexpression Cell Lysate Product Information

|                              |  |
|------------------------------|--|
| <b>Expressed Host:</b>       | HEK293 Cells   |
| <b>Products Description:</b> | Human Cell lysate that Rat EFNB1 / Ephrin-B1 transfected / overexpressed for Western blot (WB) positive control. The whole cell lysate is provided in 1X Sample Buffer (1X modified RIPA buffer+1X SDS loading buffer).        |
| <b>Sequence information:</b> | A DNA sequence encoding the rat EFNB1 (P52796) extracellular domain (Met 1-Thr 229) was expressed, fused with a polyhistidine tag at the C-terminus.   |
| <b>Predicted N Terminal:</b> | Ala 25   |
| <b>Molecule Mass:</b>        | The recombinant rat EFNB1 comprises 216 amino acids and predicts a molecular mass of 23.8 kDa. The apparent molecular mass of the ratEFNB1 is approximately 36 kDa in SDS-PAGE under reducing conditions due to glycosylation. |
| <b>Species:</b>              | Rat  |

## Ephrin-B1 Transfected / Overexpression Cell Lysate Usage Guide

|                                 |  |
|---------------------------------|--|
| <b>Preparation Method:</b>      | Cell lysate was prepared by homogenization in ice-cold modified RIPA Lysis Buffer with cocktail of protease inhibitors (Sigma). Cell debris was removed by centrifugation. Protein concentration was determined by Bradford assay (Bio-Rad protein assay, Microplate Standard assay). The cell lysate was boiled for 5 min in 1 x SDS loading buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized. |
| <b>Lysis Buffer:</b>            | Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF.   |
| <b>Quality Control Testing:</b> | 12.5% SDS-PAGE Stained with Coomassie Blue after protein purification.   |
| <b>Stability:</b>               | Samples are stable for up to twelve months from date of receipt.   |
| <b>Recommend Usage:</b>         | <ol style="list-style-type: none"><li>1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube.</li><li>2. Re-dissolve the pellet using 200µL pure water and boil for 2-5 min.</li><li>3. Store the lyophilized cell lysate at 4°C. After re-dissolution, recommend to aliquot it into smaller quantities and store at -80°C.</li></ol>  |
| <b>Storage Buffer:</b>          | 1 X Sample Buffer (1 X modified RIPA buffer+1 X SDS loading buffer).   |
| <b>Storage Instruction:</b>     | Store at 4°C. After re-dissolution, aliquot and store at -80°C.  |
| <b>Application notes:</b>       | Western blot (WB): Use at an assay dependent dilution.<br>Other Applications: Not tested.<br>Optimal dilutions/concentrations should be determined by the end user.  |

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217

● Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288

● Tel:+86-400-890-9989 ●

● <http://www.sinobiological.com>