

Rat c-MPL / CD110 / TPOR Protein (His Tag)



Sino Biological
Biological Solution Specialist

Catalog Number: 80346-R08B

General Information

Gene Name Synonym:

MPL

Protein Construction:

A DNA sequence encoding the mature form of rat MPL (Met1-Ala500) was expressed with a polyhistidine tag at the C-terminus.

Source: Rat

Expression Host: Baculovirus-Insect Cells

QC Testing

Purity: > 90 % as determined by SDS-PAGE

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Gln 22

Molecular Mass:

The recombinant rat MPL consists of 490 amino acids and predicts a molecular mass of 55.1 KDa. It migrates as an approximately 55 KDa band in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile 20mM Tris, 500mM NaCl, 3mM DTT, 10% glycerol, pH 7.5.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

CD110, also known as c-MPL, is a 635 amino acid transmembrane domain, with two extracellular cytokine receptor domains and two intracellular cytokine receptor box motifs. It is expressed at a low level in a large number of cells of hematopoietic origin. C-MPL is homologous with members of the hematopoietic receptor superfamily. Presence of anti-sense oligodeoxynucleotides of c-mpl inhibited megakaryocyte colony formation. Thrombopoietin is the ligand for c-mpl. It was shown to be the major regulator of megakaryocytopoiesis and platelet formation. Defects in c-MPL are a cause of congenital amegakaryocytic thrombocytopeniawhich is a disease characterized by isolated thrombocytopenia and megakaryocytopenia with no physical anomalies. Defects in c-MPL also cause thrombocythemia type 2 and myelofibrosis with myeloid metaplasia.

References

1. Vigon I., *et al.*, (1992), Molecular cloning and characterization of MPL, the human homolog of the v-mpl oncogene: identification of a member of the hematopoietic growth factor receptor superfamily. *Proc. Natl. Acad. Sci. U.S.A.* 89:5640-5644.
2. Mignotte V., *et al.*, (1994), Structure and transcription of the human c-mpl gene (MPL). *Genomics* 20:5-12.
3. Gregory S.G., *et al.*, (2006), The DNA sequence and biological annotation of human chromosome 1. *Nature* 441:315-321.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>