

Rat SPN / CD43 / Sialophorin Protein (Fc Tag)



Sino Biological
Biological Solution Specialist

Catalog Number: 80432-R02H

General Information

Gene Name Synonym:

SPN

Protein Construction:

A DNA sequence encoding the rat SPN (NP_001258015.1)(Met1-Gly243) was expressed with the Fc region of human IgG1 at the C-terminus.

Source: Rat

Expression Host: HEK293 Cells

QC Testing

Purity: (56.8+36.6+4.3) % as determined by SDS-PAGE

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Glu 20

Molecular Mass:

The recombinant rat SPN/Fc is a disulfide-linked homodimer. The reduced monomer comprises 465 amino acids and has a predicted molecular mass of 49.3 kDa. The apparent molecular mass of the protein is approximately 95, 57 and 50 kDa in SDS-PAGE under reducing conditions due to glycosylation.

Formulation:

Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

CD43 is an abundantly expressed molecule on the T-cell surface that shows distinct localization to the migrating T-cell uropod and the distal pole complex (DPC) opposite the immunological synapse via association with the ezrin-radixin-moesin (ERM) family of actin regulatory proteins. CD43 has a 235-amino acid (aa) extracellular domain, a 23-aa transmembrane domain, and a 123-aa cytoplasmic domain, all encoded by a single exon. The intracytoplasmic region of the protein is necessary to transduce signals; it is rich in potentially phosphorylatable threonines and serines but lacks tyrosine residues as well as catalytic activity. CD43 engagement on human peripheral blood T cells and monocytes leads to cell activation and proliferation through the generation of second messengers such as diacylglycerol and inositol phosphates, protein kinase C (PKC) activation and Ca²⁺ mobilization. In addition, CD43 ligation on human T cells induces the association of CD43 with Src family kinases, presumably through the interaction of their Src homology 3 domain with a proline-rich region of the CD43 intracytoplasmic tail. This molecule has been implicated in T cell activation, enhancing T cell response to allogeneic or mitogenic stimulation and CD43-specific signals have been reported to be sufficient to activate T cells in the absence of T cell receptor (TCR) engagement. In summary, CD43 regulates multiple T-cell functions, including T-cell activation, proliferation, apoptosis, and migration.

References

- 1.. Layseca-Espinosa E, et al. (2003) Journal of Leukocyte Biology. 74(6): 1083-93.
- 2.Cannon JL, et al. (2011) Mol Biol Cell. 22(7):954-63.
- 3.Pallant A, et al. (1989). Proc Natl Acad Sci. 86 (4): 1328-32.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>