

# Rat IL9 / IL-9 Protein (His Tag)

Catalog Number: 80459-R08B



Sino Biological  
Biological Solution Specialist

## General Information

### Gene Name Synonym:

IL9

### Protein Construction:

A DNA sequence encoding the rat IL9 (NP\_001099217.1) (Met1-Ala144) was expressed with a polyhistidine tag at the C-terminus.

Source: Rat

Expression Host: Baculovirus-Insect Cells

## QC Testing

Purity: > 90 % as determined by SDS-PAGE.

### Bio Activity:

Measured in a cell proliferation assay using MC/9-2 mouse mast cells.  
The ED<sub>50</sub> for this effect is typically 19.8-79.2 ng/mL.

### Endotoxin:

< 1.0 EU per µg protein as determined by the LAL method.

### Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Gln 19

### Molecular Mass:

The recombinant rat IL9 consists 137 amino acids and predicts a molecular mass of 15.6 kDa.

### Formulation:

Lyophilized from sterile 20 mM Tris, 500 mM NaCl, 10 % glycerol.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

## Usage Guide

### Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

### Reconstitution:

Detailed reconstitution instructions are sent along with the products.

## SDS-PAGE:



## Protein Description

Interleukin 9, also known as IL-9, is a cytokine (cell signalling molecule) belonging to the group of interleukins. IL-9 is a cytokine that acts as a regulator of a variety of hematopoietic cells. This cytokine stimulates cell proliferation and prevents apoptosis. It functions through the interleukin 9 receptor (IL-9R), which activates different signal transducer and activator (STAT) proteins and thus connects this cytokine to various biological processes. Genetic studies on a mouse model of asthma demonstrated that this cytokine is a determining factor in the pathogenesis of bronchial hyperresponsiveness. IL-9 is a key molecule that affects differentiation of TH17 cells and Treg function. IL-9 predominantly produced by TH17 cells, synergizes with TGF-β1 to differentiate naïve CD4+ T cells into TH17 cells, while IL-9 secretion by TH17 cells is regulated by IL-23. Interestingly, IL-9 enhances the suppressive functions of FoxP3+ CD4+ Treg cells in vitro, and absence of IL-9 signaling weakens the suppressive activity of nTregs in vivo, leading to an increase in effector cells and worsening of experimental autoimmune encephalomyelitis. The mechanism of IL-9 effects on TH17 and Tregs is through activation of STAT3 and STAT5 signaling. Our findings highlight a role of IL-9 as a regulator of pathogenic versus protective mechanisms of immune responses.

## References

1. Elyaman W, *et al.* (2009) IL-9 induces differentiation of TH17 cells and enhances function of FoxP3+ natural regulatory T cells. *Proc Natl Acad Sci U S A.* 106(31): 12885-90.
2. Dong Q, *et al.* (1999) IL-9 induces chemokine expression in lung epithelial cells and baseline airway eosinophilia in transgenic mice. *Eur J Immunol.* 29(7): 2130-9.
3. Kimura Y, *et al.* (1995) Sharing of the IL-2 receptor gamma chain with the functional IL-9 receptor complex. *Int Immunol.* 7(1): 115-20.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>