

Rhesus KIM-1 / TIM1 / HACVR1 Protein (His Tag)

Catalog Number: 90006-C07H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

HAVCR1

Protein Construction:

A DNA sequence encoding the rhesus HAVCR1 (BAJ61041.1) N-terminal segment (Asp 20-Val 135) was expressed with an N-terminal polyhistidine tag.

Source: Rhesus

Expression Host: HEK293 Cells

QC Testing

Purity: > 95 % as determined by SDS-PAGE

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: His

Molecular Mass:

The recombinant rhesus HAVCR1 consists of 135 amino acids and has a calculated molecular mass of 15 kDa.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

HAV cellular receptor 1 (HAVCR1), also known as Kidney injury molecule 1 (KIM-1) and T cell immunoglobulin mucin 1 (TIM-1), is a type A integral membrane glycoprotein. KIM-1 protein is widely expressed with highest levels in kidney and testis. It has been shown to play a major role as a human susceptibility gene for asthma, allergy and autoimmunity. IgA1lambda is a specific ligand of KIM-1 protein and that their association has a synergistic effect in virus-receptor interactions. KIM-1 involves in the pathogenesis of acute kidney injury. It had been confirmed that KIM-1 is a human urinary renal dysfunction biomarker. Moreover, KIM-1 protein is a novel regulatory molecule of flow-induced calcium signaling.

References

1. Tami C, *et al.* (2007) Immunoglobulin A (IgA) is a natural ligand of hepatitis A virus cellular receptor 1 (HAVCR1), and the association of IgA with HAVCR1 enhances virus-receptor interactions. *J Virol.* 81(7): 3437-46.
2. Rees AJ, *et al.* (2008) Kim-1/Tim-1: from biomarker to therapeutic target? *Nephrol Dial Transplant.* 23(11): 3394-6.
3. Chaturvedi S, *et al.* (2009) Assay validation for KIM-1: human urinary renal dysfunction biomarker. *Int J Biol Sci.* 5(2): 128-34.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>