

# Rhesus AMY2A / Alpha-amylase Protein (Fc Tag)

Catalog Number: 90012-C02H



Sino Biological  
Biological Solution Specialist

## General Information

### Gene Name Synonym:

AMY2A; AMY2B

### Protein Construction:

A DNA sequence encoding the rhesus AMY2A (H9EX43) (Met1-Leu511) was expressed with the Fc region of human IgG1 at the C-terminus.

**Source:** Rhesus

**Expression Host:** HEK293 Cells

## QC Testing

**Purity:** > 90 % as determined by SDS-PAGE

### Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

### Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

**Predicted N terminal:** Gln 16

### Molecular Mass:

The recombinant rhesus AMY2A comprises 737 amino acids and has a calculated molecular mass of 82.8 KDa.

### Formulation:

Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

## Usage Guide

### Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

**Avoid repeated freeze-thaw cycles.**

### Reconstitution:

Detailed reconstitution instructions are sent along with the products.

## SDS-PAGE:



## Protein Description

Alpha-amylase is the major form of amylase found in humans and other mammals. Amylases are secreted proteins that hydrolyze 1,4-alpha-glucoside bonds in oligosaccharides and polysaccharides, and thus catalyze the first step in digestion of dietary starch and glycogen. Alpha-amylase hydrolyses alpha bonds of large, alpha-linked polysaccharides, such as starch and glycogen, yielding glucose and maltose. Amylases is widely expressed and is most prominent in pancreatic juice and saliva, each of which has its own isoform of human  $\alpha$ -amylase. They behave differently on isoelectric focusing, and can also be separated in testing by using specific monoclonal antibodies.

## References

1.Abe A, *et al.* (2005) Complexes of Thermoactinomyces vulgaris R-47 Alpha-amylase / AMY2A 1 and pullulan model oligosaccharides provide new insight into the mechanism for recognizing substrates with alpha-(1,6) glycosidic linkages. FEBS J. 272(23):6145-53. 2.Aghajari, N, *et al.* (1998) Crystal structures of the psychrophilic Alpha-amylase / AMY2A from Alteromonas haloplantis in its native form and complexed with an inhibitor. Protein Sci. 7(3): 564-72. 3.Ramasubbu, N, *et al.* (1996) Structure of Human Salivary -Amylase at 1.6 Resolution: Implications for its Role in the Oral Cavity. Acta Crystallographica Section D Biological Crystallography. 52(3):435-46.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>