

Cynomolgus Cadherin-6 / CDH6 Protein (His Tag)



Sino Biological
Biological Solution Specialist

Catalog Number: 90910-C08H

General Information

Gene Name Synonym:

CDH6

Protein Construction:

A DNA sequence encoding the cynomolgus CDH6 (XP_005556691.1) (Met1-Gly614) was expressed with a polyhistidine tag at the C-terminus.


Source: Cynomolgus

Expression Host: HEK293 Cells

QC Testing

Purity: > 75 % as determined by SDS-PAGE.

SDS-PAGE:



Endotoxin:

<1.0 EU per µg protein as determined by the LAL method.

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Thr 22

Molecular Mass:

The recombinant cynomolgus CDH6 consists of 604 amino acids and predicts a molecular mass of 67.5 kDa.

Formulation:

Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:

Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

Protein Description

Cadherins are a family of calcium-dependent, cell-cell adhesion molecules that play an important morphoregulatory role in a wide variety of tissues. Alterations in cadherin function have been implicated in tumor progression in a number of adenocarcinomas. Cadherin-6 (CDH6), also known as K-cadherin (KCAD), is a type-II classic cadherin cell-cell adhesion molecules, which are expressed in graded or areal patterns, as well as layer-specific patterns, in the cortical plate. Human Cadherin-6 is synthesized as a 79 aa type I transmembrane glycoprotein that contains a 18 aa signal peptide, a 35 aa propeptide, a 562 aa extracellular region, a 21 aa transmembrane segment, and a 154 aa cytoplasmic domain. There are five cadherin domains of approximately 11 aa each in the extracellular region. Cadherin-6 is highly expressed in brain, cerebellum, and kidney, and may contribute to the formation of the segmental structure of the early brain, as well as the development of renal proximal tubules. Weak expression is also detected lung, pancreas, and gastric mucosa. Additionally, it is specifically expressed in the proximal tubule of normal kidneys and in renal cell cancer. Thus, Cadherin-6 is a new prognostic factor for renal cancer.

References

Paul R, et al. (1997) Cadherin-6, a cell adhesion molecule specifically expressed in the proximal renal tubule and renal cell carcinoma. *Cancer Res.* 57(13): 2741-8. Paul R, et al. (2004) Cadherin-6: a new prognostic marker for renal cell carcinoma. *J Urol.* 171(1): 97-101. Taniguchi H, et al. (2006) Classic cadherins regulate tangential migration of precerebellar neurons in the caudal hindbrain. *Development.* 133(10): 1923-31. Liu Q, et al. (2006) Cadherin-6 message expression in the nervous system of developing zebrafish. *Dev Dyn.* 235(1): 272-8.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>