

Human FCGRT & B2M Heterodimer HEK293

Cell Lysate (WB positive control)



Sino Biological Inc.
Biological Solution Specialist

Catalog Number: CT071-H27HL

FCGRT & B2M Heterodimer Transfected / Overexpression Cell Lysate Product Information

Expressed Host:	HEK293 Cells
Products Description:	Human Cells that Human FCGRT & B2M Heterodimer Heterodimer transfected / overexpressed for Western blot (WB) positive control. The whole cell lysate is provided in 1X Sample Buffer (1X modified RIPA buffer+1X SDS loading buffer).
Sequence information:	A DNA sequence encoding the extracellular domain (Met1-Ser297) of human FCGRT (NP_004098.1) was fused with a C-terminal polyhistidine tagged AVI tag at the C-terminus, constructed the plasmid 1; A DNA sequence encoding the human B2M (NP_004039.1) (Met1-Met119) constructed the plasmid 2. The two plasmids were co-expressed and the human FCGRT/B2M heterodimer was purified.
Predicted N Terminal:	Ala 24 & Ile 21
Molecule Mass:	The recombinant heterodimer of human FCGRT/B2M comprises 399 (300+99) amino acids and has a calculated molecular mass of 45.3 (33.6+11.7) KDa.
Species:	Human

FCGRT & B2M Heterodimer Transfected / Overexpression Cell Lysate Usage Guide

Preparation Method:	Cell lysate was prepared by homogenization in ice-cold modified RIPA Lysis Buffer with cocktail of protease inhibitors (Sigma). Cell debris was removed by centrifugation. Protein concentration was determined by Bradford assay (Bio-Rad protein assay, Microplate Standard assay). The cell lysate was boiled for 5 min in 1 x SDS loading buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis Buffer:	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF.
Quality Control Testing:	12.5% SDS-PAGE Stained with Coomassie Blue after protein purification.
Stability:	Samples are stable for up to twelve months from date of receipt.
Recommend Usage:	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boil for 2-5 min. 3. Store the lyophilized cell lysate at 4°C. After re-dissolution, recommend to aliquot it into smaller quantities and store at -80°C.
Storage Buffer:	1 X Sample Buffer (1 X modified RIPA buffer+1 X SDS loading buffer).
Storage Instruction:	Store at 4°C. After re-dissolution, aliquot and store at -80°C.
Application notes:	Western blot (WB): Use at an assay dependent dilution. Other Applications: Not tested. Optimal dilutions/concentrations should be determined by the end user.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 ● Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 ● Tel:+86-400-890-9989 ● <http://www.sinobiological.com>